



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12P 21/06, C12N 1/20, 15/00, C07H 21/04, C07K 1/00, 14/52	A1	(11) International Publication Number: WO 98/07880 (43) International Publication Date: 26 February 1998 (26.02.98)
(21) International Application Number: PCT/US96/13282 (22) International Filing Date: 16 August 1996 (16.08.96) (71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): YU, Guo-Liang [CN/US]; 13524 Straw Bale Lane, Darnestown, MD 20878 (US). NI, Jian [CN/US]; 5502 Manorfield Road, Rockville, MD 20853 (US). ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). (74) Agents: GOLDSTEIN, Jorge, A. et al.; Sterne, Kessler, Goldstein & Fox P.L.L.C., Suite 600, 1100 New York Avenue, N.W., Washington, DC 20005-3934 (US).		(81) Designated States: AM, AU, BG, BR, BY, CA, CN, CZ, EE, FI, GE, HU, IL, JP, KG, KP, KR, KZ, LT, LV, MD, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, UA, US, UZ, VN, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: HUMAN ENDOKINE ALPHA (57) Abstract The present invention concerns a novel member of the tumor necrosis factor (TNF) family of cytokines. In particular, isolated nucleic acid molecules are provided encoding the endokine alpha protein. Endokine alpha polypeptides are also provided, as are vectors, host cells and recombinant methods for producing the same. Also provided are diagnostic and therapeutic methods concerning TNF family-related disorders.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Human Endokine Alpha

Background of the Invention

Field of the Invention

5 The present invention relates to an endokine alpha protein. In particular, isolated nucleic acid molecules are provided encoding the endokine alpha protein. Endokine alpha polypeptides are also provided, as are vectors, host cells and recombinant methods for producing the same.

Related Art

10 The cytokine known as tumor necrosis factor- α (TNF α ; also termed cachectin) is a protein secreted primarily by monocytes and macrophages in response to endotoxin or other stimuli as a soluble homotrimer of 17 kD protein subunits (Smith, R.A. *et al.*, *J. Biol. Chem.* 262:6951-6954 (1987)). A membrane-bound 26 kD precursor form of TNF has also been described (Kriegler, M. *et al.*, *Cell* 53:45-53 (1988)).

15 Accumulating evidence indicates that TNF is a regulatory cytokine with pleiotropic biological activities. These activities include: inhibition of lipoprotein lipase synthesis ("cachectin" activity) (Beutler, B. *et al.*, *Nature* 316:552 (1985)), activation of polymorphonuclear leukocytes (Klebanoff, S.J. *et al.*, *J. Immunol.* 136:4220 (1986); Perussia, B., *et al.*, *J. Immunol.* 138:765 (1987)), inhibition of
20 cell growth or stimulation of cell growth (Vilcek, J. *et al.*, *J. Exp. Med.* 163:632 (1986); Sugarman, B.J. *et al.*, *Science* 230:943 (1985); Lachman, L.B. *et al.*, *J. Immunol.* 138:2913 (1987)), cytotoxic action on certain transformed cell types (Lachman, L.B. *et al.*, *supra*; Darzynkiewicz, Z. *et al.*, *Canc. Res.* 44:83 (1984)), antiviral activity (Kohase, M. *et al.*, *Cell* 45:659 (1986); Wong, G.H.W. *et al.*,
25 *Nature* 323:819 (1986)), stimulation of bone resorption (Bertolini, D.R. *et al.*, *Nature* 319:516 (1986); Saklatvala, J., *Nature* 322:547 (1986)), stimulation of collagenase and prostaglandin E2 production (Dayer, J.-M. *et al.*, *J. Exp. Med.*

-2-

162:2163 (1985)); and immunoregulatory actions, including activation of T cells (Yokota, S. *et al.*, *J. Immunol.* 140:531 (1988)), B cells (Kehrl, J.H. *et al.*, *J. Exp. Med.* 166:786 (1987)), monocytes (Philip, R. *et al.*, *Nature* 323:86 (1986)), thymocytes (Ranges, G.E. *et al.*, *J. Exp. Med.* 167:1472 (1988)), and stimulation of the cell-surface expression of major histocompatibility complex (MHC) class I and class II molecules (Collins, T. *et al.*, *Proc. Natl. Acad. Sci. USA* 83:446 (1986); Pujol-Borrel, R. *et al.*, *Nature* 326:304 (1987)).

TNF is noted for its pro-inflammatory actions which result in tissue injury, such as induction of procoagulant activity on vascular endothelial cells (Pober, J.S. *et al.*, *J. Immunol.* 136:1680 (1986)), increased adherence of neutrophils and lymphocytes (Pober, J.S. *et al.*, *J. Immunol.* 138:3319 (1987)), and stimulation of the release of platelet activating factor from macrophages, neutrophils and vascular endothelial cells (Camussi, G. *et al.*, *J. Exp. Med.* 166:1390 (1987)).

Recent evidence implicates TNF in the pathogenesis of many infections (Cerami, A. *et al.*, *Immunol. Today* 9:28 (1988)), immune disorders, neoplastic pathology, e.g., in cachexia accompanying some malignancies (Oloff, A. *et al.*, *Cell* 50:555 (1987)), and in autoimmune pathologies and graft-versus host pathology (Piguet, P.-F. *et al.*, *J. Exp. Med.* 166:1280 (1987)). The association of TNF with cancer and infectious pathologies is often related to the host's catabolic state. A major problem in cancer patients is weight loss, usually associated with anorexia. The extensive wasting which results is known as "cachexia" (Kern, K.A. *et al.*, *J. Parent. Enter. Nutr.* 12:286-298 (1988)). Cachexia includes progressive weight loss, anorexia, and persistent erosion of body mass in response to a malignant growth. The cachectic state is thus associated with significant morbidity and is responsible for the majority of cancer mortality. A number of studies have suggested that TNF is an important mediator of the cachexia in cancer, infectious pathology, and in other catabolic states.

TNF is thought to play a central role in the pathophysiological consequences of Gram-negative sepsis and endotoxic shock (Michie, H.R. *et al.*, *Br. J. Surg.* 76:670-671 (1989); Debets, J.M.H. *et al.*, *Second Vienna Shock*

-3-

Forum, p.463-466 (1989); Simpson, S.Q. *et al.*, *Crit. Care Clin.* 5:27-47 (1989)), including fever, malaise, anorexia, and cachexia. Endotoxin is a potent monocyte/macrophage activator which stimulates production and secretion of TNF (Kornbluth, S.K. *et al.*, *J. Immunol.* 137:2585-2591 (1986)) and other cytokines. Because TNF could mimic many biological effects of endotoxin, it was concluded to be a central mediator responsible for the clinical manifestations of endotoxin-related illness. TNF and other monocyte-derived cytokines mediate the metabolic and neurohormonal responses to endotoxin (Michie, H.R. *et al.*, *N. Eng. J. Med.* 318:1481-1486 (1988)). Endotoxin administration to human volunteers produces acute illness with flu-like symptoms including fever, tachycardia, increased metabolic rate and stress hormone release (Revhaug, A. *et al.*, *Arch. Surg.* 123:162-170 (1988)). Elevated levels of circulating TNF have also been found in patients suffering from Gram-negative sepsis (Waage, A. *et al.*, *Lancet* 1:355-357 (1987); Hammerle, A.F. *et al.*, *Second Vienna Shock Forum* p. 715-718 (1989); Debets, J.M.H. *et al.*, *Crit. Care Med.* 17:489-497 (1989); Calandra, T. *et al.*, *J. Infec. Dis.* 161:982-987 (1990)).

Passive immunotherapy directed at neutralizing TNF may have a beneficial effect in Gram-negative sepsis and endotoxemia, based on the increased TNF production and elevated TNF levels in these pathology states, as discussed above.

Antibodies to a "modulator" material which was characterized as cachectin (later found to be identical to TNF) were disclosed by Cerami *et al.* (EPO Patent Publication 0,212,489, March 4, 1987). Such antibodies were said to be useful in diagnostic immunoassays and in therapy of shock in bacterial infections. Rubin *et al.* (EPO Patent Publication 0,218,868, April 22, 1987) disclosed monoclonal antibodies to human TNF, the hybridomas secreting such antibodies, methods of producing such antibodies, and the use of such antibodies in immunoassay of TNF. Yone *et al.* (EPO Patent Publication 0,288,088, October 26, 1988) disclosed anti-TNF antibodies, including mAbs, and their utility in immunoassay diagnosis of pathologies, in particular Kawasaki's pathology and bacterial infection. The body fluids of patients with Kawasaki's

-4-

pathology (infantile acute febrile mucocutaneous lymph node syndrome; Kawasaki, T., *Allergy* 16:178 (1967); Kawasaki, T., *Shonica (Pediatrics)* 26:935 (1985)) were said to contain elevated TNF levels which were related to progress of the pathology (Yone *et al.*, *supra*).

5 Other investigators have described mAbs specific for recombinant human TNF which had neutralizing activity in vitro (Liang, C-M. *et al. Biochem. Biophys. Res. Comm.* 137:847-854 (1986); Meager, A. *et al., Hybridoma* 6:305-311 (1987); Fendly *et al., Hybridoma* 6:359-369 (1987); Bringman, T.S. *et al., Hybridoma* 6:489-507 (1987); Hirai, M. *et al., J. Immunol. Meth.* 96:57-62 (1987); Moller, A. *et al. (Cytokine* 2:162-169 (1990)). Some of these mAbs were used to map epitopes of human TNF and develop enzyme immunoassays (Fendly *et al., supra*; Hirai *et al., supra*; Moller *et al., supra*) and to assist in the purification of recombinant TNF (Bringman *et al., supra*). However, these studies do not provide a basis for producing TNF neutralizing antibodies that can be used for in vivo diagnostic or therapeutic uses in humans, due to immunogenicity, lack of specificity and/or pharmaceutical suitability.

15 Neutralizing antisera or mAbs to TNF have been shown in mammals other than man to abrogate adverse physiological changes and prevent death after lethal challenge in experimental endotoxemia and bacteremia. This effect has been demonstrated, e.g., in rodent lethality assays and in primate pathology model systems (Mathison, J.C. *et al., J. Clin. Invest.* 81:1925-1937 (1988); Beutler, B. *et al., Science* 229:869-871 (1985); Tracey, K.J. *et al., Nature* 330:662-664 (1987); Shimamoto, Y. *et al., Immunol. Lett.* 17:311-318 (1988); Silva, A.T. *et al., J. Infect. Dis.* 162:421-427 (1990); Opal, S.M. *et al., J. Infect. Dis.* 161:1148-1152 (1990); Hinshaw, L.B. *et al., Circ. Shock* 30:279-292 (1990)).

20 To date, experience with anti-TNF mAb therapy in humans has been limited but shows beneficial therapeutic results, e.g., in arthritis and sepsis. See, e.g., Elliott, M.J. *et al., Baillieres Clin. Rheumatol.* 9:633-52 (1995); Feldmann M, *et al., Ann. N. Y. Acad. Sci. USA* 766:272-8 (1995); van der Poll, T. *et al., Shock* 3:1-12 (1995); Wherry *et al., Crit. Care Med.* 21:S436-40 (1993); Tracey K.J., *et al., Crit. Care Med.* 21:S415-22 (1993).

-5-

Sequence analysis of cytokine receptors has defined several subfamilies of membrane proteins (1) the Ig superfamily, (2) the hematopoietin (cytokine receptor superfamily and (3) the tumor necrosis factor (TNF)/nerve growth factor (NGF) receptor superfamily (for review of TNF superfamily see, Gruss and Dower, *Blood* 85(12):3378-3404 (1995) and Aggarwal and Natarajan, *Eur. Cytokine Netw.*, 7(2):93-124 (1996)). The TNF/NGF receptor superfamily contains at least 10 different proteins. Gruss and Dower, *supra*. Ligands for these receptors have been identified and belong to at least two cytokine superfamilies. Gruss and Dower, *supra*.

Accordingly, there is a need to provide cytokines similar to TNF that are involved in pathological conditions. Such novel cytokines could be used to make novel antibodies or other antagonists that bind these TNF-like cytokines for therapy of TNF-like disorders.

Summary of the Invention

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding a cytokine that is similar to TNF and is believed to have similar biological effects and activities. This cytokine is named endokine alpha, and includes endokine alpha polypeptides having at least a portion of the amino acid sequence in FIG. 1 (SEQ ID NO:2) or an amino acid sequence encoded by the cDNA clone deposited in a bacterial host as ATCC Deposit Number 97640 on June 27, 1996. The nucleotide sequence, which was determined by sequencing the deposited endokine alpha cDNA clone, contains an open reading frame encoding a polypeptide of about 169 amino acid residues including an N-terminal methionine, an intracellular domain of about 17 amino acid residues, a transmembrane domain of about 26 amino acids, an extracellular domain of about 126 amino acids, and a deduced molecular weight for the complete protein of about 19 kDa. The 126 amino acid sequence of the expected mature endokine alpha protein is shown in FIG. 1 (SEQ ID NO:2) (residues 44 to 169).

-6-

In another aspect, the invention provides isolated nucleic acid molecules encoding the endokine alpha polypeptide having an amino acid sequence encoded by the cDNA of the clone deposited as ATCC Deposit No. 97640 on June 27, 1996. Preferably, the nucleic acid molecule will encode the mature polypeptide encoded by the above-described deposited cDNA.

The invention is further directed to nucleic acid fragments of the nucleic acid molecules described herein. Preferred nucleic acid fragments include nucleic acid molecules which encode: a polypeptide comprising the endokine alpha intracellular domain (amino acid residues from about 1 to about 17 in FIG. 1 (SEQ ID NO:2)); a polypeptide comprising the endokine alpha transmembrane domain (amino acid residues from about 18 to about 43 in FIG. 1 (SEQ ID NO:2)); and a polypeptide comprising the endokine alpha extracellular domain (amino acid residues from about 44 to about 169 in FIG. 1 (SEQ ID NO:2)).

Further embodiments of the invention include an isolated nucleic acid molecule having a nucleotide sequence that is at least 90% identical, and more preferably at least 95%, 96%, 97%, 98%, or 99% identical to the nucleotide sequence of any of the nucleic acid molecules describe herein.

The present invention also relates to recombinant vectors which include the isolated nucleic acid molecules of the present invention, host cells containing the recombinant vectors, and the production of endokine alpha polypeptides by recombinant techniques.

The polypeptides of the present invention include the polypeptide encoded by the deposited cDNA, the polypeptide of FIG. 1 (SEQ ID NO:2) (in particular the mature polypeptide), as well as polypeptides having an amino acid sequence with at least 90% similarity, more preferably at least 95% similarity to the amino acid sequence of the polypeptide encoded by the deposited cDNA, the polypeptide of FIG. 1 (SEQ ID NO:2), or a fragment thereof. Further polypeptides of the present invention include polypeptides having an amino acid sequence at least 80% identical, more preferably, at least 90% or 95% identical to the amino acid sequence of the polypeptide encoded by the deposited cDNA, the polypeptide of FIG. 1 (SEQ ID NO:2), or a fragment thereof.

-7-

Preferred polypeptide fragments according to the present invention include a polypeptide comprising: the endokine alpha intracellular domain, the endokine alpha transmembrane domain, and the endokine alpha extracellular domain.

5 The invention further provides methods for isolating antibodies that bind specifically to an endokine alpha polypeptide having an amino acid sequence as described above. Such antibodies may be useful diagnostically or therapeutically as antagonists in the treatment of endokine alpha- and/or TNF-related disorders.

Brief Description of the Figures

10 FIG. 1 shows the nucleotide (SEQ ID NO:1) and deduced amino acid (SEQ ID NO:2) sequences of the endokine alpha protein. Amino acids 1 to 17 represent the intracellular domain, amino acids 18 to 43 the transmembrane domain (the underlined sequence), and amino acids 44 to 169 the extracellular domain (the remaining sequence).

15 FIG. 2 shows the regions of similarity between the amino acid sequences of the endokine alpha protein (SEQ ID NO:2), tissue necrosis factor α (TNF- α) (SEQ ID NO:3), and TNF- β (SEQ ID NO:4). The J. Hein method was used with PAM 250 residue weight table. Shading with solid black indicates residues that match consensus exactly.

20 FIG. 3 provides an analysis of the endokine alpha amino acid sequence. Alpha, beta, turn and coil regions; hydrophilicity and hydrophobicity; amphipathic regions; flexible regions; antigenic index and surface probability are shown. In the "Antigenic Index - Jameson-Wolf" graph, amino acid residues 44-54, 57-68, 69-78, 94-105, 108-132 and 148-158 in FIG. 1 correspond to the
25 shown highly antigenic regions of the endokine alpha protein.

Detailed Description of the Invention

The present invention provides isolated nucleic acid molecules comprising a polynucleotide encoding an endokine alpha protein having an amino acid sequence shown in FIG. 1 (SEQ ID NO:2), which was determined by sequencing a cloned cDNA. Endokine alpha is a novel member of the tumor necrosis factor (TNF) ligand family and shares sequence homology with human TNF α and related TNF family members (FIG. 2). The nucleotide sequence shown in FIG. 1 (SEQ ID NO:1) was obtained by sequencing a cDNA clone, which was deposited on June 27, 1996, at the American Type Culture Collection, 12301 Park Lawn Drive, Rockville, Maryland 20852, and given accession number 97640. The deposited clone is contained in the pBluescript SK(-) plasmid (Stratagene, LaJolla, CA).

Nucleic Acid Molecules

Unless otherwise indicated, all nucleotide sequences determined by sequencing a DNA molecule herein were determined using an automated DNA sequencer (such as the Model 373 from Applied Biosystems, Inc.), and all amino acid sequences of polypeptides encoded by DNA molecules determined herein were predicted by translation of a DNA sequence determined as above. Therefore, as is known in the art for any DNA sequence determined by this automated approach, any nucleotide sequence determined herein may contain some errors. Nucleotide sequences determined by automation are typically at least about 90% identical, more typically at least about 95% to at least about 99.99% identical to the actual nucleotide sequence of the sequenced DNA molecule. The actual sequence can be more precisely determined by other approaches including manual DNA sequencing methods well known in the art. As is also known in the art, a single insertion or deletion in a determined nucleotide sequence compared to the actual sequence will cause a frame shift in translation of the nucleotide sequence such that the expected amino acid sequence

encoded by a determined nucleotide sequence will be completely different from the amino acid sequence actually encoded by the sequenced DNA molecule, beginning at the point of such an insertion or deletion.

5 Unless otherwise indicated, each "nucleotide sequence" set forth herein is presented as a sequence of deoxyribonucleotides (abbreviated A, G, C and T). However, by "nucleotide sequence" of a nucleic acid molecule or polynucleotide is intended, for a DNA molecule or polynucleotide, a sequence of deoxyribonucleotides, and for an RNA molecule or polynucleotide, the corresponding sequence of ribonucleotides (A, G, C and U) where each thymidine
10 deoxynucleotide (T) in the specified deoxynucleotide sequence is replaced by the ribonucleotide uridine (U). For instance, reference to an RNA molecule having the sequence of FIG. 1 (SEQ ID NO:1) set forth using deoxyribonucleotide abbreviations is intended to indicate an RNA molecule having a sequence in which each deoxynucleotide A, G or C of SEQ ID NO:1
15 has been replaced by the corresponding ribonucleotide A, G or C, and each deoxynucleotide T has been replaced by a ribonucleotide U.

Using the information provided herein, such as the nucleotide sequence in FIG. 1, a nucleic acid molecule of the present invention encoding an endokine alpha polypeptide can be obtained using standard cloning and screening
20 procedures, such as those for cloning cDNAs using mRNA as starting material. Illustrative of the invention, the nucleic acid molecule described in FIG. 1 (SEQ ID NO:1) was discovered in a cDNA library derived from human brain striatum. Expressed sequence tags corresponding to a portion of the endokine alpha cDNA were also found in several endothelial libraries and a fetal liver library.

25 The endokine alpha gene contains an open reading frame encoding a protein of about 169 amino acid residues, an intracellular domain of about 17 amino acids (amino acid residues from about 1 to about 17 in FIG. 1 (SEQ ID NO:2)), a transmembrane domain of about 26 amino acids (amino acid residues from about 18 to about 43 in FIG. 1 (SEQ ID NO:2)), an extracellular domain of
30 about 126 amino acids (amino acid residues from about 44 to about 169 in FIG. 1 (SEQ ID NO:2)); and a deduced molecular weight of about 19 kDa. The

-10-

endokine alpha protein shown in FIG. 1 (SEQ ID NO: 2) is about 30% similar and about 22% identical to human TNF- α , which can be accessed on GenBank as Accession No. U42764.

5 As one of ordinary skill would appreciate, due to the possibilities of sequencing errors discussed above, as well as the variability of cleavage sites for leaders in different known proteins, the actual endokine alpha polypeptide encoded by the deposited cDNA comprises about 169 amino acids, but can be anywhere in the range of about 154-184 amino acids. It will also be appreciated by reasonable persons of skill in the art that, depending on the criteria used, the exact 'address' of the above-described endokine alpha protein domains may differ. 10 Thus, for example, the exact location of the endokine alpha intracellular, transmembrane and extracellular domains shown in FIG. 1 (SEQ ID NO:2) may vary slightly (e.g., the exact address may differ by about 1 to about 5 residues compared to that shown in FIG. 1) depending on the criteria used to define the domain. 15

As indicated, nucleic acid molecules of the present invention may be in the form of RNA, such as mRNA, or in the form of DNA, including, for instance, cDNA and genomic DNA obtained by cloning or produced synthetically. The DNA may be double-stranded or single-stranded. Single-stranded DNA can be 20 the coding strand, also known as the sense strand, or it can be the non-coding strand, also referred to as the anti-sense strand.

By "isolated" nucleic acid molecule(s) is intended a nucleic acid molecule, DNA or RNA, which has been removed from its native environment. For example, recombinant DNA molecules contained in a vector are considered 25 isolated for the purposes of the present invention. Further examples of isolated DNA molecules include recombinant DNA molecules maintained in heterologous host cells or purified (partially or substantially) DNA molecules in solution. Isolated RNA molecules include *in vivo* or *in vitro* RNA transcripts of the DNA molecules of the present invention. Isolated nucleic acid molecules according to 30 the present invention further include such molecules produced synthetically.

-11-

Isolated nucleic acid molecules of the present invention include DNA molecules comprising the open reading frame (ORF) shown in FIG. 1 (SEQ ID NO:1) and further include nucleic acid molecules substantially different than all or part of the ORF sequence shown in FIG. 1 (SEQ ID NO:1) but which, due to the degeneracy of the genetic code, still encode the endokine alpha protein or a fragment thereof. Of course, the genetic code is well known in the art. Thus, it would be routine for one skilled in the art to generate the degenerate variants described above.

In another aspect, the invention provides isolated nucleic acid molecules encoding the endokine alpha polypeptide having an amino acid sequence encoded by the cDNA of the clone deposited as ATCC Deposit No. 97640 on June 27, 1996. Preferably, the nucleic acid molecule will encode the mature polypeptide encoded by the above-described deposited cDNA clone. The invention further provides an isolated nucleic acid molecule having the nucleotide sequence shown in FIG. 1 (SEQ ID NO:1) or the nucleotide sequence of the endokine alpha cDNA contained in the above-described deposited clone, or a nucleic acid molecule having a sequence complementary to one of the above sequences. Such isolated molecules, particularly DNA molecules, are useful as probes for gene mapping by *in situ* hybridization with chromosomes and for detecting expression of the endokine alpha gene in human tissue, for instance, by Northern blot analysis. As described in detail below, detecting altered endokine alpha gene expression in certain tissues or bodily fluids is indicative of certain disorders.

The present invention is further directed to fragments of the isolated nucleic acid molecules described herein. By a fragment of an isolated nucleic acid molecule having the nucleotide sequence of the deposited cDNA or the nucleotide sequence shown in FIG. 1 (SEQ ID NO. 1) is intended fragments at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt in length which are useful as diagnostic probes and primers as discussed herein. Of course, larger fragments 50-1500 nt in length are also useful according to the present invention as are fragments corresponding to most, if not all, of the nucleotide sequence of

-12-

the deposited cDNA or as shown in FIG. 1 (SEQ ID NO. 1) . By a fragment at least 20 nt in length, for example, is intended fragments which include 20 or more contiguous bases from the nucleotide sequence of the deposited cDNA or the nucleotide sequence as shown in FIG. 1 (SEQ ID NO. 1). Since the gene has been deposited and the nucleotide sequence shown in FIG. 1 (SEQ ID NO 1) is provided, generating such DNA fragments would be routine to the skilled artisan. For example, restriction endonuclease cleavage or shearing by sonication could easily be used to generate fragments of various sizes. Alternatively, such fragments could be generated synthetically.

Preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising the endokine alpha intracellular domain (amino acid residues from about 1 to about 17 in FIG. 1 (SEQ ID NO 2)); a polypeptide comprising the endokine alpha transmembrane domain (amino acid residues from about 18 to about 43 in FIG. 1 (SEQ ID NO 2)); and a polypeptide comprising the endokine alpha extracellular domain (amino acid residues from about 44 to about 169 in FIG. 1 (SEQ ID NO. 2)).

Further preferred nucleic acid fragments of the present invention include nucleic acid molecules encoding epitope-bearing portions of the endokine alpha protein. In particular, such nucleic acid fragments of the present invention include nucleic acid molecules encoding: a polypeptide comprising amino acid residues from about 44 to about 158 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 44 to about 54 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 57 to about 68 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 69 to about 78 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 94 to about 105 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 108 to about 132 in FIG. 1 (SEQ ID NO:2); and a polypeptide comprising amino acid residues from about 148 to about 158 in FIG. 1 (SEQ ID NO:2). The inventors have determined that the above polypeptide fragments are antigenic regions of the endokine alpha protein.

-13-

Methods for determining other such epitope-bearing portions of the endokine alpha protein are described in detail below.

In another aspect, the invention provides an isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a portion of the polynucleotide in a nucleic acid molecule of the invention described above, for instance, the cDNA clone contained in ATCC Deposit 97640 made on June 27, 1996. By "stringent hybridization conditions" is intended overnight incubation at 42°C in a solution comprising: 50% formamide, 5x SSC (150 mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65°C. By a polynucleotide which hybridizes to a "portion" of a polynucleotide is intended a polynucleotide (either DNA or RNA) hybridizing to at least about 15 nucleotides (nt), and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably about 30-70 nt of the reference polynucleotide. These are useful as diagnostic probes and primers as discussed above and in more detail below.

Of course, polynucleotides hybridizing to a larger portion of the reference polynucleotide (*e.g.*, the deposited cDNA clone), for instance, a portion 50-500 nt in length, or even to the entire length of the reference polynucleotide, are also useful as probes according to the present invention, as are polynucleotides corresponding to most, if not all, of the nucleotide sequence of the deposited cDNA or the nucleotide sequence as shown in FIG. 1 (SEQ ID NO:1). By a portion of a polynucleotide of "at least 20 nt in length," for example, is intended 20 or more contiguous nucleotides from the nucleotide sequence of the reference polynucleotide, (*e.g.*, the deposited cDNA or the nucleotide sequence as shown in FIG. 1 (SEQ ID NO:1)). As indicated, such portions are useful diagnostically either as a probe according to conventional DNA hybridization techniques or as primers for amplification of a target sequence by the polymerase chain reaction (PCR), as described, for instance, in Sambrook, J. *et al.*, eds., *Molecular Cloning, A Laboratory Manual*, 2nd. edition, Cold Spring Harbor Laboratory Press, Cold

-14-

Spring Harbor, NY (1989), the entire disclosure of which is hereby incorporated herein by reference.

Since an endokine alpha cDNA clone has been deposited and its nucleotide sequence is provided in FIG. 1 (SEQ ID NO:1), generating polynucleotides which hybridize to a portion of the endokine alpha cDNA molecule would be routine to the skilled artisan. For example, restriction endonuclease cleavage or shearing by sonication of the endokine alpha cDNA clone could easily be used to generate DNA portions of various sizes which are polynucleotides that hybridize to a portion of the endokine alpha cDNA molecule. Alternatively, the hybridizing polynucleotides of the present invention could be generated synthetically according to known techniques.

Of course, a polynucleotide which hybridizes only to a poly A sequence (such as the 3' terminal poly(A) tract of the endokine alpha cDNA shown in FIG. 1 (SEQ ID NO:1)), or to a complementary stretch of T (or U) residues, would not be included in a polynucleotide of the invention used to hybridize to a portion of a nucleic acid of the invention, since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically any double-stranded cDNA clone).

As indicated, nucleic acid molecules of the present invention that encode an endokine alpha protein may include, but are not limited to, those encoding the amino acid sequence of the mature polypeptide, by itself; the coding sequence for the mature polypeptide and additional sequences, such as a pre-, or pro- or prepro- protein sequence; the coding sequence of the mature polypeptide, with or without the aforementioned additional coding sequences, together with additional, non-coding sequences, including for example, but not limited to introns and non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription, mRNA processing - including splicing and polyadenylation signals, e.g., ribosome binding and stability of mRNA; an additional coding sequence which codes for additional amino acids, such as those which provide additional functionalities. Thus, for instance, the sequence encoding the polypeptide can be fused to a marker sequence, such as a sequence

-15-

encoding a peptide which facilitates purification of the fused polypeptide. In certain preferred embodiments of this aspect of the invention, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (Qiagen, Inc.), among others, many of which are publicly and/or commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. The "HA" tag is another peptide useful for purification which corresponds to an epitope derived from the influenza hemagglutinin (HA) protein, which has been described by Wilson *et al.*, *Cell* 37:767 (1984). Other such fusion proteins include the endokine alpha protein fused to Fc at the N- or C-terminus.

The present invention further relates to variants of the nucleic acid molecules of the present invention, which encode portions, analogs or derivatives of the endokine alpha protein. Variants can occur naturally, such as a natural allelic variant. By an "allelic variant" is intended one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. Non-naturally occurring variants can be produced, *e.g.*, using art-known mutagenesis techniques.

Such variants include those produced by nucleotide substitutions, deletions or additions. The substitutions, deletions or additions may involve one or more nucleotides. The variants can be altered in coding or non-coding regions or both. Alterations in the coding regions can produce conservative or non-conservative amino acid substitutions, deletions or additions. Especially preferred among these are silent substitutions, additions and deletions, which do not alter the properties and activities of the endokine alpha protein or portions thereof. Also especially preferred in this regard are conservative substitutions. Most highly preferred are nucleic acid molecules encoding the mature endokine alpha protein having the amino acid sequence shown in FIG. 1 (SEQ ID NO:2) or the mature endokine alpha amino acid sequence encoded by the deposited cDNA clone.

-16-

Further embodiments of the invention include isolated nucleic acid molecules comprising a polynucleotide having a nucleotide sequence at least 90% identical, and more preferably at least 95%, 96%, 97%, 98%, or 99% identical to (a) a nucleotide sequence encoding the full-length endokine alpha polypeptide having the complete amino acid sequence shown in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640; (b) a nucleotide sequence encoding the endokine alpha extracellular domain having the amino acid sequence at positions from about 44 to about 169 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640; (c) a nucleotide sequence encoding the endokine alpha transmembrane domain having the amino acid sequence at positions from about 18 to about 43 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640; (d) a nucleotide sequence encoding the endokine alpha intracellular domain having the amino acid sequence at positions from about 1 to about 17 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640; or (e) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c) or (d).

By a polynucleotide having a nucleotide sequence at least, for example, 95% "identical" to a reference nucleotide sequence encoding an endokine alpha polypeptide is intended that the nucleotide sequence of the polynucleotide is identical to the reference sequence except that the polynucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the endokine alpha polypeptide. In other words, to obtain a polynucleotide having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. These mutations of the reference sequence may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually

-17-

among nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular nucleic acid molecule is at least 90, 95%, 96%, 97%, 98%, or 99% identical to, for instance, the nucleotide sequence shown in FIG. 1 or to the nucleotides sequence of the deposited cDNA clone can be determined conventionally using known computer programs such as the BESTFIT program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711). BESTFIT uses the local homology algorithm of Smith and Waterman, *Adv. Appl. Math.* 2:482-489 (1981), to find the best segment of homology between two sequences. When using BESTFIT or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference nucleotide sequence and that gaps in homology of up to 5% of the total number of nucleotides in the reference sequence are allowed.

The present application is directed to such nucleic acid molecules which are at least 90%, 95%, 96%, 97%, 98%, or 99% identical to a nucleic acid sequence described above irrespective of whether they encode a polypeptide having endokine alpha protein activity. This is because, even where a particular nucleic acid molecule does not encode a polypeptide having endokine alpha activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having endokine alpha activity include, *inter alia*, (1) isolating the endokine alpha gene or allelic variants thereof from a cDNA library; (2) *in situ* hybridization (FISH) to metaphase chromosomal spreads to provide precise chromosomal location of the endokine alpha gene as described in Verma *et al.*, *Human Chromosomes: a Manual of Basic Techniques*, Pergamon

-18-

Press, New York (1988); and (3) Northern Blot analysis for detecting endokine alpha mRNA expression in specific tissues.

5 Preferred, however, are such nucleic acid molecules having sequences at least 90%, 95%, 96%, 97%, 98%, or 99% identical to a nucleic acid sequence described above which do, in fact, encode a polypeptide having endokine alpha protein activity. By "a polypeptide having endokine alpha activity" is intended polypeptides exhibiting similar, but not necessarily identical, activity as compared to the endokine alpha protein as measured in a particular biological assay. Endokine alpha activity can be assayed according to known methods. For
10 example, a cytotoxicity assay or cell proliferation assay can be used where endokine alpha polypeptides are added to cells in culture and the effect of the endokine on the cells is determined by measuring the decrease or increase in cell numbers.

Of course, due to the degeneracy of the genetic code, one of ordinary skill
15 in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 90%, 95%, 96%, 97%, 98%, or 99% identical to a nucleic acid sequence described above will encode a polypeptide "having endokine alpha protein activity." In fact, since degenerate variants all encode the same polypeptide, this will be clear to the skilled artisan even without
20 performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having endokine alpha protein activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein
25 function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid).

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie, J.U. *et al.*, *Science* 247:1306-1310 (1990), wherein the authors indicate that there are two main approaches for
30 studying the tolerance of an amino acid sequence to change. The first method relies on the process of evolution, in which mutations are either accepted or

-19-

rejected by natural selection. The second approach uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene and selections or screens to identify sequences that maintain functionality. As the authors state, these studies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at a certain position of the protein. For example, most buried amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Other such phenotypically silent substitutions are described in Bowie, J.U., *et al.*, *supra*, and the references cited therein.

Vectors and Host Cells

The present invention also relates to vectors which include the isolated DNA molecules of the present invention, host cells which are genetically engineered with the recombinant vectors, and the production of endokine alpha polypeptides or portions thereof by recombinant techniques.

Recombinant constructs may be introduced into host cells using well known techniques such as infection, transduction, transfection, transvection, electroporation and transformation. The vector may be, for example, a phage, plasmid, viral or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

The polynucleotides may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged *in vitro* using an appropriate packaging cell line and then transduced into host cells.

Preferred are vectors comprising cis-acting control regions to the polynucleotide of interest. Appropriate trans-acting factors may be supplied by

-20-

the host, supplied by a complementing vector or supplied by the vector itself upon introduction into the host.

5 In certain preferred embodiments in this regard, the vectors provide for specific expression, which may be inducible and/or cell type-specific. Particularly preferred among such vectors are those inducible by environmental factors that are easy to manipulate, such as temperature and nutrient additives.

10 Expression vectors useful in the present invention include chromosomal-, episomal- and virus-derived vectors, e.g., vectors derived from bacterial plasmids, bacteriophage, yeast episomes, yeast chromosomal elements, viruses such as baculoviruses, papova viruses, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as cosmids and phagemids. See, e.g., Ausubel, *infra*; Sambrook, *infra*.

15 The DNA insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the *E. coli*: *lac*, *trp* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination and, in the transcribed region, a ribosome binding site for translation.
20 The coding portion of the mature transcripts expressed by the constructs will include a translation initiating AUG at the beginning and a termination codon appropriately positioned at the end of the polypeptide to be translated.

25 As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase or neomycin resistance for eukaryotic cell culture and tetracycline or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells; insect cells such as *Drosophila* S2 and *Spodoptera Sf9* cells; animal cells such as CHO, COS and Bowes melanoma cells; and plant cells. Appropriate culture media and
30 conditions for the above-described host cells are known in the art.

-21-

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from Qiagen; pBS vectors, Phagescript vectors, Bluescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene; and ptrc99a, pKK223-3, pKK233-3, pDR540, pRIT5 available from Pharmacia. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Other suitable vectors will be readily apparent to the skilled artisan.

Among known bacterial promoters suitable for use in the present invention include the *E. coli lacI* and *lacZ* promoters, the T3 and T7 promoters, the *gpt* promoter, the lambda PR and PL promoters and the *trp* promoter. Suitable eukaryotic promoters include the CMV immediate early promoter, the HSV thymidine kinase promoter, the early and late SV40 promoters, the promoters of retroviral LTRs, such as those of the Rous sarcoma virus (RSV), and metallothionein promoters, such as the mouse metallothionein-I promoter.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods in Molecular Biology* (1986).

Transcription of the DNA encoding the polypeptides of the present invention by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp that act to increase transcriptional activity of a promoter in a given host cell-type. Examples of enhancers include the SV40 enhancer, which is located on the late side of the replication origin at bp 100 to 270, the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers.

For secretion of the translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide.

-22-

The signals may be endogenous to the polypeptide or they may be heterologous signals.

The polypeptide may be expressed in a modified form, such as a fusion protein, and may include not only secretion signals but also additional heterologous functional regions. In a further example, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence in the host cell, during purification or during subsequent handling and storage. Also, as indicated, a region(s) also may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to polypeptides to engender secretion or excretion, to improve stability and to facilitate purification, among others, are familiar and routine techniques in the art. A preferred fusion protein comprises a heterologous region from immunoglobulin that is useful to solubilize receptors. For example, EP A 0,464,533 (*also*, Canadian counterpart 2,045,869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in the fusion protein is thoroughly advantageous for use in therapy and diagnosis and thus results, for example, in improved pharmacokinetic properties (EP A 0,232,262). On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified in the advantageous manner described. This is the case when Fc portion proves to be a hindrance to use in therapy and diagnosis, for example when the fusion protein is to be used as antigen for immunizations. In drug discovery, for example, human proteins, such as, hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists (for example, hIL-5). See, D. Bennett *et al.*, *Journal of Molecular Recognition* 8:52-58 (1995) and K. Johanson *et al.*, *The Journal of Biological Chemistry* 270(16):9459-9471 (1995).

The endokine alpha protein can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate

-23-

or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention include naturally purified products, products of chemical synthetic procedures, and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes.

Endokine Alpha Polypeptides and Peptides

The invention further provides an isolated endokine alpha polypeptide having the amino acid sequence encoded by the deposited cDNA, or the amino acid sequence in FIG. 1 (SEQ ID NO:2), or a peptide or polypeptide comprising a portion of the above polypeptides. The terms "peptide" and "oligopeptide" are considered synonymous (as is commonly recognized) and each term can be used interchangeably as the context requires to indicate a chain of at least two amino acids coupled by peptidyl linkages. The word "polypeptide" is used herein for chains containing more than ten amino acid residues. All oligopeptide and polypeptide formulas or sequences herein are written from left to right and in the direction from amino terminus to carboxy terminus.

By "isolated" polypeptide or protein is intended a polypeptide or protein removed from its native environment. For example, recombinantly produced polypeptides and proteins expressed in recombinant host cells are considered isolated for purposes of the invention as are native or recombinant polypeptides and proteins which have been substantially purified by any suitable technique

-24-

such as, for example, the one-step method described in Smith and Johnson, *Gene* 67:31-40 (1988).

5 It will be recognized in the art that some amino acid sequence of the endokine alpha polypeptide can be varied without significant effect of the structure or function of the protein. If such differences in sequence are contemplated, it should be remembered that there will be critical areas on the protein which determine activity. In general, it is possible to replace residues which form the tertiary structure, provided that residues performing a similar function are used. In other instances, the type of residue may be completely
10 unimportant if the alteration occurs at a non-critical region of the protein.

Thus, the invention further includes variations of the endokine alpha polypeptide which show substantial endokine alpha polypeptide activity or which include regions of endokine alpha protein such as the protein fragments discussed below. Such mutants include deletions, insertions, inversions, repeats, and type
15 substitutions (for example, substituting one hydrophilic residue for another, but not strongly hydrophilic for strongly hydrophobic as a rule). Small changes or such "neutral" amino acid substitutions will generally have little effect on activity.

Typically seen as conservative substitutions are the replacements, one for
20 another, among the aliphatic amino acids Ala, Val, Leu and Ile; interchange of the hydroxyl residues Ser and Thr, exchange of the acidic residues Asp and Glu, substitution between the amide residues Asn and Gln, exchange of the basic residues Lys and Arg and replacements among the aromatic residues Phe, Tyr.

As indicated in detail above, further guidance concerning which amino
25 acid changes are likely to be phenotypically silent (i.e., are not likely to have a significant deleterious effect on a function) can be found in Bowie, J.U., *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

The polypeptides of the present invention include polypeptides having an
30 amino acid sequence as shown in FIG. 1 (SEQ ID NO:2) or as encoded by the deposited cDNA such as: the full-length protein, the intracellular domain, the

-25-

transmembrane domain, and the extracellular domain, as well as polypeptides which have at least 90% similarity, more preferably at least 95% similarity, and still more preferably at least 96%, 97%, 98% or 99% similarity to a polypeptide described herein. Further polypeptides of the present invention include polypeptides at least 80% identical, more preferably at least 90% or 95% identical, still more preferably at least 96%, 97%, 98% or 99% identical to a polypeptide described herein, and also include portions of such polypeptides with at least 30 amino acids and more preferably at least 50 amino acids.

By "% similarity" for two polypeptides is intended a similarity score produced by comparing the amino acid sequences of the two polypeptides using the BESTFIT program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711) and the default settings for determining similarity. BESTFIT uses the local homology algorithm of Smith and Waterman, *Adv. Appl. Math.* 2:482-489 (1981), to find the best segment of similarity between two sequences.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of an endokine alpha polypeptide is intended that the amino acid sequence of the polypeptide is identical to the reference sequence except that the polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the reference amino acid sequence of the endokine alpha polypeptide. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence may be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence may be inserted into the reference sequence. These alterations of the reference sequence may occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

-26-

As a practical matter, whether any particular polypeptide is at least 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence shown in FIG. 1 (SEQ ID NO:2) or to the amino acid sequence encoded by deposited cDNA clone can be determined conventionally using known computer programs such the BESTFIT program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI 53711. When using BESTFIT or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

As described in detail below, the polypeptides of the present invention can be used to raise polyclonal and monoclonal antibodies, which are useful in diagnostic assays for detecting endokine alpha protein expression as described below or as agonists and antagonists capable of inhibiting endokine alpha protein function. Further, such polypeptides can be used in the yeast two-hybrid system to "capture" endokine alpha protein binding proteins which are also candidate agonist and antagonist according to the present invention. The yeast two hybrid system is described in Fields and Song, *Nature* 340:245-246 (1989).

In another aspect, the invention provides a peptide or polypeptide comprising an epitope-bearing portion of a polypeptide of the invention. The epitope of this polypeptide portion is an immunogenic or antigenic epitope of a polypeptide of the invention. An "immunogenic epitope" is defined as a part of a protein that elicits an antibody response when the whole protein is the immunogen. These immunogenic epitopes are believed to be confined to a few loci on the molecule. On the other hand, a region of a protein molecule to which an antibody can bind is defined as an "antigenic epitope." The number of immunogenic epitopes of a protein generally is less than the number of antigenic

-27-

epitopes. See, for instance, Geysen, H.M. *et al.*, *Proc. Natl. Acad. Sci. USA* 81:3998-4002 (1984).

As to the selection of peptides or polypeptides bearing an antigenic epitope (i.e., that contain a region of a protein molecule to which an antibody can bind), it is well known in that art that relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, for instance, Sutcliffe, J.G. *et al.*, *Science* 219:660-666 (1983). Peptides capable of eliciting protein-reactive sera are frequently represented in the primary sequence of a protein, can be characterized by a set of simple chemical rules, and are confined neither to immunodominant regions of intact proteins (i.e., immunogenic epitopes) nor to the amino or carboxyl terminals. Peptides that are extremely hydrophobic and those of six or fewer residues generally are ineffective at inducing antibodies that bind to the mimicked protein; longer, soluble peptides, especially those containing proline residues, usually are effective. Sutcliffe *et al.*, *supra*, at 661. For instance, 18 of 30 peptides designed according to these guidelines, containing 8-39 residues covering 75% of the sequence of the influenza virus hemagglutinin HA1 polypeptide chain, induced antibodies that reacted with the HA1 protein or intact virus; and 12/12 peptides from the MuLV polymerase and 18/18 from the rabies glycoprotein induced antibodies that precipitated the respective proteins.

Antigenic epitope-bearing peptides and polypeptides of the invention are therefore useful to raise antibodies, including monoclonal antibodies, that bind specifically to a polypeptide of the invention. Thus, a high proportion of hybridomas obtained by fusion of spleen cells from donors immunized with an antigen epitope-bearing peptide generally secrete antibody reactive with the native protein. Sutcliffe *et al.*, *supra*, at 663. The antibodies raised by antigenic epitope-bearing peptides or polypeptides are useful to detect the mimicked protein, and antibodies to different peptides may be used for tracking the fate of various regions of a protein precursor which undergoes posttranslational processing. The peptides and anti-peptide antibodies may be used in a variety of qualitative or quantitative assays for the mimicked protein, for instance in

-28-

competition assays since it has been shown that even short peptides (e.g., about 9 amino acids) can bind and displace the larger peptides in immunoprecipitation assays. See, for instance, Wilson, I.A. *et al.*, *Cell* 37:767-778 (1984) at 777. The anti-peptide antibodies of the invention also are useful for purification of the mimicked protein, for instance, by adsorption chromatography using methods well known in the art.

Antigenic epitope-bearing peptides and polypeptides of the invention designed according to the above guidelines preferably contain a sequence of at least seven, more preferably at least nine and most preferably between about 15 to about 30 amino acids contained within the amino acid sequence of a polypeptide of the invention. However, peptides or polypeptides comprising a larger portion of an amino acid sequence of a polypeptide of the invention, containing about 30 to about 50 amino acids, or any length up to and including the entire amino acid sequence of a polypeptide of the invention, also are considered epitope-bearing peptides or polypeptides of the invention and also are useful for inducing antibodies that react with the mimicked protein. Preferably, the amino acid sequence of the epitope-bearing peptide is selected to provide substantial solubility in aqueous solvents (i.e., the sequence includes relatively hydrophilic residues and highly hydrophobic sequences are preferably avoided); and sequences containing proline residues are particularly preferred.

Non-limiting examples of antigenic polypeptides that can be used to generate endokine-specific polyclonal and monoclonal antibodies include: a polypeptide comprising amino acid residues from about 44 to about 158 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 44 to about 54 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 57 to about 68 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 69 to about 78 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 94 to about 105 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 108 to about 132 in FIG. 1 (SEQ ID NO:2); and a polypeptide comprising amino acid residues from about 148 to about 158 in FIG. 1 (SEQ ID NO:2). As

-29-

indicated above, the inventors have determined that the above polypeptide fragments are antigenic regions of the endokine alpha protein.

The epitope-bearing peptides and polypeptides of the invention may be produced by any conventional means for making peptides or polypeptides including recombinant means using nucleic acid molecules of the invention. For instance, a short epitope-bearing amino acid sequence may be fused to a larger polypeptide which acts as a carrier during recombinant production and purification, as well as during immunization to produce anti-peptide antibodies. Epitope-bearing peptides also may be synthesized using known methods of chemical synthesis. For instance, Houghten has described a simple method for synthesis of large numbers of peptides, such as 10-20 mg of 248 different 13 residue peptides representing single amino acid variants of a segment of the HA1 polypeptide which were prepared and characterized (by ELISA-type binding studies) in less than four weeks. *See, Houghten, R.A., Proc. Natl. Acad. Sci. USA* 82:5131-5135 (1985). This "Simultaneous Multiple Peptide Synthesis (SMPS)" process is further described in U.S. Patent No. 4,631,211 to Houghten *et al.* (1986). In this procedure the individual resins for the solid-phase synthesis of various peptides are contained in separate solvent-permeable packets, enabling the optimal use of the many identical repetitive steps involved in solid-phase methods. A completely manual procedure allows 500-1000 or more syntheses to be conducted simultaneously. Houghten *et al., supra*, at 5134.

Epitope-bearing peptides and polypeptides of the invention are used to induce antibodies according to methods well known in the art. *See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow, M. et al., Proc. Natl. Acad. Sci. USA* 82:910-914; and Bittle, F.J. *et al., J. Gen. Virol.* 66:2347-2354 (1985). Generally, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling of the peptide to a macromolecular carrier, such as keyhole limpet hemocyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine may be coupled to carrier using a linker such as m-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS), while other

-30-

peptides may be coupled to carrier using a more general linking agent such as glutaraldehyde.

5 Animals such as rabbits, rats and mice are immunized with either free or carrier-coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg peptide or carrier protein and Freund's adjuvant. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an
10 immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

 Immunogenic epitope-bearing peptides of the invention, i.e., those parts of a protein that elicit an antibody response when the whole protein is the
15 immunogen, are identified according to methods known in the art. For instance, Geysen *et al.* (1984), *supra*, discloses a procedure for rapid concurrent synthesis on solid supports of hundreds of peptides of sufficient purity to react in an enzyme-linked immunosorbent assay. Interaction of synthesized peptides with antibodies is then easily detected without removing them from the support. In
20 this manner a peptide bearing an immunogenic epitope of a desired protein may be identified routinely by one of ordinary skill in the art.

 For instance, the immunologically important epitope in the coat protein of foot-and-mouth disease virus was located by Geysen *et al.* with a resolution of seven amino acids by synthesis of an overlapping set of all 208 possible
25 hexapeptides covering the entire 213 amino acid sequence of the protein. Then, a complete replacement set of peptides in which all 20 amino acids were substituted in turn at every position within the epitope were synthesized, and the particular amino acids conferring specificity for the reaction with antibody were determined. Thus, peptide analogs of the epitope-bearing peptides of the
30 invention can be made routinely by this method. U.S. Patent No. 4,708,781 to

-31-

Geysen (1987) further describes this method of identifying a peptide bearing an immunogenic epitope of a desired protein.

Further still, U.S. Patent No. 5,194,392 to Geysen (1990) describes a general method of detecting or determining the sequence of monomers (amino acids or other compounds) which is a topological equivalent of the epitope (i.e., a "mimotope") which is complementary to a particular paratope (antigen binding site) of an antibody of interest. More generally, U.S. Patent No. 4,433,092 to Geysen (1989) describes a method of detecting or determining a sequence of monomers which is a topographical equivalent of a ligand which is complementary to the ligand binding site of a particular receptor of interest. Similarly, U.S. Patent No. 5,480,971 to Houghten, R. A. *et al.* (1996) on Peralkylated Oligopeptide Mixtures discloses linear C₁-C₇-alkyl peralkylated oligopeptides and sets and libraries of such peptides, as well as methods for using such oligopeptide sets and libraries for determining the sequence of a peralkylated oligopeptide that preferentially binds to an acceptor molecule of interest. Thus, non-peptide analogs of the epitope-bearing peptides of the invention also can be made routinely by these methods.

The present inventors have discovered that the endokine alpha protein is a 169 residue protein exhibiting three main structural domains. The intracellular domain was identified within residues from about 1 to about 17 in FIG. 1 (SEQ ID NO:2). The transmembrane domain was identified within residues from about 18 to about 43 in FIG. 1 (SEQ ID NO:2). The extracellular domain was identified within residues from about 44 to about 169 in FIG. 1 (SEQ ID NO:2). Thus, the invention further provides preferred endokine alpha protein fragments comprising a polypeptide selected from: the endokine alpha intracellular domain, the transmembrane domain and the endokine alpha extracellular domain.

The extracellular domain of the endokine alpha protein can be combined with parts of the constant domain of immunoglobulins (IgG), resulting in chimeric polypeptides. These fusion proteins show an increased half-life *in vivo*. This has been shown, e.g., for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant

-32-

regions of the heavy or light chains of mammalian immunoglobulins (EP A 394,827; Traunecker *et al.*, *Nature* 331:84-86 (1988)). Fusion proteins that have a disulfide-linked dimeric structure due to the IgG part can also be more efficient in binding and neutralizing the ligands than the monomeric extracellular domains alone (Fountoulakis *et al.*, *J. Biochem.* 270:3958-3964 (1995)).

The entire disclosure of each document cited in this section on "Polypeptides and Peptides" is hereby incorporated herein by reference.

Endokine Alpha Related Disorder Diagnosis

Endokine alpha is a new member of the TNF family of cytokines. For endokine alpha related disorders, it is believed that substantially altered (increased or decreased) levels of endokine alpha gene expression can be detected in tissue or other cells or bodily fluids (e.g., sera, plasma, urine, synovial fluid or spinal fluid) taken from an individual having such a disorder, relative to a "standard" endokine alpha gene expression level, that is, the endokine alpha expression level in tissue or bodily fluids from an individual not having the disorder. Thus, the invention provides a diagnostic method useful during diagnosis of an endokine alpha-related disorder, which involves measuring the expression level of the gene encoding the endokine alpha protein in tissue or other cells or body fluid from an individual and comparing the measured gene expression level with a standard endokine alpha gene expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of an endokine alpha related disorder.

By individual is intended mammalian individuals, preferably humans. By "measuring the expression level of the gene encoding the endokine alpha protein" is intended qualitatively or quantitatively measuring or estimating the level of the endokine alpha protein or the level of the mRNA encoding the endokine alpha protein in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the endokine alpha protein level or mRNA level in a second

-33-

biological sample). Preferably, the endokine alpha protein level or mRNA level in the first biological sample is measured or estimated and compared to a standard endokine alpha protein level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the disorder or being determined by averaging levels from a population of individuals not having a disorder involving endokine alpha. As will be appreciated in the art, once a standard endokine alpha protein level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source which contains endokine alpha protein or mRNA. As indicated, biological samples include body fluids (such as sera, plasma, urine, synovial fluid and spinal fluid) which contain secreted mature endokine alpha protein, or tissue sources found to express endokine alpha. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the preferred source.

The present invention is useful for diagnosis of various endokine alpha - related disorders in mammals, preferably humans, as similar to TNF-like disorders known in the art or as presented herein. These include disorders associated with immunomodulation and inflammation, cell proliferation, angiogenesis, tumor metastases, apoptosis, sepsis and endotoxemia.

Total cellular RNA can be isolated from a biological sample using any suitable technique such as the single-step guanidinium-thiocyanate-phenol-chloroform method described in Chomczynski and Sacchi, *Anal. Biochem.* 162:156-159 (1987). Levels of mRNA encoding an endokine alpha polypeptide are then assayed using any appropriate method. These include Northern blot analysis, S1 nuclease mapping, the polymerase chain reaction (PCR), reverse transcription in combination with the polymerase chain reaction (RT-PCR), and reverse transcription in combination with the ligase chain reaction (RT-LCR).

Northern blot analysis can be performed as described in Harada *et al.*, *Cell* 63:303-312 (1990). Briefly, total RNA is prepared from a biological sample as

-34-

described above. For the Northern blot, the RNA is denatured in an appropriate buffer (such as glyoxal/dimethyl sulfoxide/sodium phosphate buffer), subjected to agarose gel electrophoresis, and transferred onto a nitrocellulose filter. After the RNAs have been linked to the filter by a UV linker, the filter is prehybridized
5 in a solution containing formamide, SSC, Denhardt's solution, denatured salmon sperm, SDS, and sodium phosphate buffer. Endokine alpha protein cDNA labeled according to any appropriate method (such as the ³²P-multiprimed DNA labeling system (Amersham)) is used as probe. After hybridization overnight, the filter is washed and exposed to x-ray film. cDNA for use as probe according to
10 the present invention is described in the sections above and will preferably at least 15 bp in length.

S1 mapping can be performed as described in Fujita *et al.*, *Cell* 49:357-367 (1987). To prepare probe DNA for use in S1 mapping, the sense strand of above-described cDNA is used as a template to synthesize labeled antisense
15 DNA. The antisense DNA can then be digested using an appropriate restriction endonuclease to generate further DNA probes of a desired length. Such antisense probes are useful for visualizing protected bands corresponding to the target mRNA (*i.e.*, mRNA encoding the endokine alpha protein). Northern blot analysis can be performed as described above.

Preferably, levels of mRNA encoding the endokine alpha protein are assayed using the RT-PCR method described in Makino *et al.*, *Technique* 2:295-301 (1990). By this method, the radioactivities of the "amplicons" in the polyacrylamide gel bands are linearly related to the initial concentration of the target mRNA. Briefly, this method involves adding total RNA isolated from a
20 biological sample in a reaction mixture containing a RT primer and appropriate buffer. After incubating for primer annealing, the mixture can be supplemented with a RT buffer, dNTPs, DTT, RNase inhibitor and reverse transcriptase. After incubation to achieve reverse transcription of the RNA, the RT products are then subject to PCR using labeled primers. Alternatively, rather than labeling the
25 primers, a labeled dNTP can be included in the PCR reaction mixture. PCR amplification can be performed in a DNA thermal cycler according to
30

-35-

conventional techniques. After a suitable number of rounds to achieve amplification, the PCR reaction mixture is electrophoresed on a polyacrylamide gel. After drying the gel, the radioactivity of the appropriate bands (corresponding to the mRNA encoding the endokine alpha protein) is quantified using an imaging analyzer. RT and PCR reaction ingredients and conditions, reagent and gel concentrations, and labeling methods are well known in the art. Variations on the RT-PCR method will be apparent to the skilled artisan.

Any set of oligonucleotide primers which will amplify reverse transcribed target mRNA can be used and can be designed as described in the sections above.

Assaying endokine alpha protein levels in a biological sample can occur using any art-known method. Preferred for assaying endokine alpha protein levels in a biological sample are antibody-based techniques. For example, endokine alpha protein expression in tissues can be studied with classical immunohistological methods. In these, the specific recognition is provided by the primary antibody (polyclonal or monoclonal) but the secondary detection system can utilize fluorescent, enzyme, or other conjugated secondary antibodies. As a result, an immunohistological staining of tissue section for pathological examination is obtained. Tissues can also be extracted, e.g., with urea and neutral detergent, for the liberation of endokine alpha protein for Western-blot or dot/slot assay (Jalkanen, M., *et al.*, *J. Cell. Biol.* 101:976-985 (1985); Jalkanen, M., *et al.*, *J. Cell. Biol.* 105:3087-3096 (1987)). In this technique, which is based on the use of cationic solid phases, quantitation of endokine alpha protein can be accomplished using isolated endokine alpha protein as a standard. This technique can also be applied to body fluids. With these samples, a molar concentration of endokine alpha protein will aid to set standard values of endokine alpha protein content for different body fluids, like serum, plasma, urine, synovial fluid, spinal fluid, etc. The normal appearance of endokine alpha protein amounts can then be set using values from healthy individuals, which can be compared to those obtained from a test subject.

Other antibody-based methods useful for detecting endokine alpha protein levels include immunoassays, such as the enzyme linked immunosorbent assay

-36-

(ELISA) and the radioimmunoassay (RIA). For example, endokine alpha protein-specific monoclonal antibodies can be used both as an immunoadsorbent and as an enzyme-labeled probe to detect and quantify the endokine alpha protein. The amount of endokine alpha protein present in the sample can be calculated by reference to the amount present in a standard preparation using a linear regression computer algorithm. Such an ELISA for detecting a tumor antigen is described in Iacobelli *et al.*, *Breast Cancer Research and Treatment* 11:19-30 (1988). In another ELISA assay, two distinct specific monoclonal antibodies can be used to detect endokine alpha protein in a body fluid. In this assay, one of the antibodies is used as the immunoadsorbent and the other as the enzyme-labeled probe.

The above techniques may be conducted essentially as a "one-step" or "two-step" assay. The "one-step" assay involves contacting endokine alpha protein with immobilized antibody and, without washing, contacting the mixture with the labeled antibody. The "two-step" assay involves washing before contacting the mixture with the labeled antibody. Other conventional methods may also be employed as suitable. It is usually desirable to immobilize one component of the assay system on a support, thereby allowing other components of the system to be brought into contact with the component and readily removed from the sample.

Suitable enzyme labels include, for example, those from the oxidase group, which catalyze the production of hydrogen peroxide by reacting with substrate. Glucose oxidase is particularly preferred as it has good stability and its substrate (glucose) is readily available. Activity of an oxidase label may be assayed by measuring the concentration of hydrogen peroxide formed by the enzyme-labeled antibody/substrate reaction. Besides enzymes, other suitable labels include radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium ($^{99\text{m}}\text{Tc}$), and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying endokine alpha protein levels in a biological sample obtained from an individual, endokine alpha protein can also be detected *in vivo* by imaging. Antibody labels or markers for *in vivo* imaging of endokine

-37-

alpha protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

A endokine alpha protein-specific antibody or antibody portion which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{111}In , $^{99\text{m}}\text{Tc}$), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously or intraperitoneally) into the mammal to be examined for a disorder. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moieties needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody portion will then preferentially accumulate at the location of cells which contain endokine alpha protein. *In vivo* tumor imaging is described in S. W. Burchiel *et al.*, "Immunopharmacokinetics of Radiolabeled Antibodies and Their Portions" (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, Burchiel, S.W. and Rhodes, B.A. eds., Masson Publishing Inc. (1982)).

Endokine alpha-protein specific antibodies for use in the present invention can be raised against the intact endokine alpha protein or an antigenic polypeptide portion thereof, which may presented together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse) or, if it is long enough (at least about 25 amino acids), without a carrier.

As used herein, the term "antibody" (Ab) or "monoclonal antibody" (Mab) is meant to include intact molecules as well as antibody portions (such as, for example, Fab and F(ab')_2 portions) which are capable of specifically binding to endokine alpha protein. Fab and F(ab')_2 portions lack the Fc portion of intact antibody, clear more rapidly from the circulation, and may have less non-specific

-38-

tissue binding of an intact antibody (Wahl *et al.*, *J. Nucl. Med.* 24:316-325 (1983)). Thus, these portions are preferred.

5 The antibodies of the present invention may be prepared by any of a variety of methods. For example, cells expressing the endokine alpha protein or an antigenic portion thereof can be administered to an animal in order to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of endokine alpha protein is prepared and purified as described above to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

10 In the most preferred method, the antibodies of the present invention are monoclonal antibodies (or endokine alpha protein binding portions thereof). Such monoclonal antibodies can be prepared using hybridoma technology (*see, e.g.*, Colligan, *Current Protocols in Immunology*, Wiley Interscience, New York (1990-1996); Harlow & Lane, *Antibodies: A Laboratory Manual*, Chs. 6-9, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1988); Ausubel, *infra*, at Chapter 11, these references entirely incorporated herein by reference).

15 In general, such procedures involve immunizing an animal (preferably a mouse) with an endokine alpha polypeptide antigen or with an endokine alpha polypeptide-expressing cell. Suitable cells can be recognized by their capacity to bind anti-endokine alpha protein antibody. Such cells may be cultured in any suitable tissue culture medium (*e.g.*, Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), supplemented with about 10 µg/l of nonessential amino acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin). The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention (*e.g.*, parent myeloma cell line (SP₂O), available from the American Type Culture Collection (ATCC) (Rockville, Maryland, USA)). After fusion, the resulting hybridoma cells are selectively maintained in HAT medium, and then cloned by limiting dilution as described by Wands *et al.*, *Gastroenterology* 80:225-232

-39-

(1981); Harlow & Lane, *infra*, Chapter 7. The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the endokine alpha antigen.

Alternatively, additional antibodies capable of binding to the endokine alpha protein antigen may be produced in a two-step procedure through the use of anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, endokine alpha protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma cells are screened to identify clones which produce an antibody whose ability to bind to the endokine alpha protein-specific antibody can be blocked by the endokine alpha protein antigen. Such antibodies comprise anti-idiotypic antibodies to the endokine alpha protein-specific antibody and can be used to immunize an animal to induce formation of further endokine alpha protein-specific antibodies.

It will be appreciated that Fab and $F(ab')_2$ and other portions of the antibodies of the present invention may be used according to the methods disclosed herein. Such portions are typically produced by proteolytic cleavage, using enzymes such as papain (to produce Fab portions) or pepsin (to produce $F(ab')_2$ portions). Alternatively, endokine alpha protein-binding portions can be produced through the application of recombinant DNA technology or through synthetic chemistry.

Where *in vivo* imaging is used to detect enhanced levels of endokine alpha protein for diagnosis in humans, it may be preferable to use "humanized" chimeric monoclonal antibodies. Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric antibodies are known in the art. See, for review, Morrison, *Science* 229:1202 (1985); Oi *et al.*, *BioTechniques* 4:214 (1986); Cabilly *et al.*, U.S. Patent No. 4,816,567; Taniguchi *et al.*, EP 171496; Morrison *et al.*, EP 173494; Neuberger *et al.*, WO 8601533; Robinson

-40-

et al., WO 8702671; Boulianne *et al.*, *Nature* 312:643 (1984); Neuberger *et al.*, *Nature* 314:268 (1985).

Further suitable labels for the endokine alpha protein-specific antibodies of the present invention are provided below. Examples of suitable enzyme labels
5 include malate dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast-alcohol dehydrogenase, alpha-glycerol phosphate dehydrogenase, triose phosphate isomerase, peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, beta-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, and acetylcholine esterase.

10 Examples of suitable radioisotopic labels include ^3H , ^{111}In , ^{125}I , ^{131}I , ^{32}P , ^{35}S , ^{14}C , ^{51}Cr , ^{57}To , ^{58}Co , ^{59}Fe , ^{75}Se , ^{152}Eu , ^{90}Y , ^{67}Cu , ^{217}Ci , ^{211}At , ^{212}Pb , ^{47}Sc , ^{109}Pd , etc. ^{111}In and $^{99\text{m}}\text{Tc}$ are preferred isotopes where *in vivo* imaging is used since its avoids the problem of dehalogenation of the ^{125}I or ^{131}I -labeled monoclonal antibody by the liver. In addition, this radionucleotide has a more favorable
15 gamma emission energy for imaging (Perkins *et al.*, *Eur. J. Nucl. Med.* 10:296-301 (1985); Carasquillo *et al.*, *J. Nucl. Med.* 28:281-287 (1987)). For example, ^{111}In coupled to monoclonal antibodies with 1-(p-isothiocyanatobenzyl)-DPTA has shown little uptake in non-tumorous tissues, particularly the liver, and therefore enhances specificity of tumor localization (Esteban *et al.*, *J. Nucl. Med.*
20 28:861-870 (1987)).

Examples of suitable non-radioactive isotopic labels include ^{157}Gd , ^{55}Mn , ^{162}Dy , ^{52}Tr , and ^{56}Fe .

Examples of suitable fluorescent labels include an ^{152}Eu label, a fluorescein label, an isothiocyanate label, a rhodamine label, a phycoerythrin
25 label, a phycocyanin label, an allophycocyanin label, an o-phthaldehyde label, and a fluorescamine label.

Examples of suitable toxin labels include diphtheria toxin, ricin, and cholera toxin.

30 Examples of chemiluminescent labels include a luminal label, an isoluminal label, an aromatic acridinium ester label, an imidazole label, an

-41-

acridinium salt label, an oxalate ester label, a luciferin label, a luciferase label, and an aequorin label.

Examples of nuclear magnetic resonance contrasting agents include heavy metal nuclei such as Gd, Mn, and Fe.

5 Typical techniques for binding the above-described labels to antibodies are provided by Kennedy *et al.* (*Clin. Chim. Acta* 70:1-31 (1976)), and Schurs *et al.* (*Clin. Chim. Acta* 81:1-40 (1977)). Coupling techniques mentioned in the latter are the glutaraldehyde method, the periodate method, the dimaleimide method, the m-maleimidobenzyl-N-hydroxy-succinimide ester method, all of
10 which methods are incorporated by reference herein.

Chromosome Assays

The nucleic acid molecules of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to and can hybridize with a particular location on an individual human chromosome.
15 Moreover, there is a current need for identifying particular sites on the chromosome. Few chromosome marking reagents based on actual sequence data (repeat polymorphisms) are presently available for marking chromosomal location. The mapping of DNAs to chromosomes according to the present invention is an important first step in correlating those sequences with genes
20 associated with disease.

In certain preferred embodiments in this regard, the cDNA herein disclosed is used to clone genomic DNA of an endokine alpha protein gene. This can be accomplished using a variety of well known techniques and libraries, which generally are available commercially. The genomic DNA then is used for
25 *in situ* chromosome mapping using well known techniques for this purpose. Typically, in accordance with routine procedures for chromosome mapping, some trial and error may be necessary to identify a genomic probe that gives a good *in situ* hybridization signal.

-42-

In some cases, in addition, sequences can be mapped to chromosomes by preparing PCR primers (preferably 15-25 bp) from the cDNA. Computer analysis of the 3' untranslated region of the gene is used to rapidly select primers that do not span more than one exon in the genomic DNA, thus complicating the amplification process. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to the primer will yield an amplified portion.

PCR mapping of somatic cell hybrids is a rapid procedure for assigning a particular DNA to a particular chromosome. Using the present invention with the same oligonucleotide primers, sublocalization can be achieved with panels of portions from specific chromosomes or pools of large genomic clones in an analogous manner. Other mapping strategies that can similarly be used to map to its chromosome include in situ hybridization, prescreening with labeled flow-sorted chromosomes and preselection by hybridization to construct chromosome specific-cDNA libraries.

Fluorescence *in situ* hybridization ("FISH") of a cDNA clone to a metaphase chromosomal spread can be used to provide a precise chromosomal location in one step. This technique can be used with probes from the cDNA as short as 50 or 60 bp. For a review of this technique, see Verma *et al.*, *Human Chromosomes: A Manual of Basic Techniques*, Pergamon Press, New York (1988).

Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found, for example, in V. McKusick, *Mendelian Inheritance in Man*, available on-line through Johns Hopkins University, Welch Medical Library. The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes).

Next, it is necessary to determine the differences in the cDNA or genomic sequence between affected and unaffected individuals. If a mutation is observed

-43-

in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease.

With current resolution of physical mapping and genetic mapping techniques, a cDNA precisely localized to a chromosomal region associated with the disease could be one of between 50 and 500 potential causative genes. (This assumes 1 megabase mapping resolution and one gene per 20 kb).

Endokine Alpha Protein and Antibody Therapy

As indicated above, TNF is noted for its pro-inflammatory actions which result in tissue injury, such as induction of procoagulant activity on vascular endothelial cells (Pober, J.S. *et al.*, *J. Immunol.* 136:1680 (1986)), increased adherence of neutrophils and lymphocytes (Pober, J.S. *et al.*, *J. Immunol.* 138:3319 (1987)), and stimulation of the release of platelet activating factor from macrophages, neutrophils and vascular endothelial cells (Camussi, G. *et al.*, *J. Exp. Med.* 166:1390 (1987)). Recent evidence implicates TNF in the pathogenesis of many infections (Cerami, A. *et al.*, *Immunol. Today* 9:28 (1988)), immune disorders, neoplastic pathology, e.g., in cachexia accompanying some malignancies (Oloff, A. *et al.*, *Cell* 50:555 (1987)), and in autoimmune pathologies and graft-versus host pathology (Piguet, P.-F. *et al.*, *J. Exp. Med.* 166:1280 (1987)). A number of studies have suggested that TNF is an important mediator of the cachexia in cancer, infectious pathology, and in other catabolic states.

Thus, the endokine alpha protein of the present invention can be used for tumor targeting, preferably, after conjugation with radioisotopes or cytostatic drugs (Gruss and Dower, *Blood* 85(12):3378-3404 (1995)). Endokine alpha can be used in patients with melanoma and sarcoma for tumor regression and extension of patient life span through a local injection or used in isolated limb perfusion (Aggarwal and Natarajan, *Eur. Cytokine Netw.* 7(2):92-124 (1996)).

The endokine alpha of the present invention can also have a therapeutic role in specific situations, for example, activity against viral, bacterial, yeast,

fungal, and other infections (including toxoplasma gondii, schistosoma mansoni, listeria monocytogens and BCG). These effects of endokine alpha can be indirect and thus preferably, mediated through activation of macrophages, eosinophils, fibroblasts, or neutrophils.

5 TNF is also thought to play a central role in the pathophysiological consequences of Gram-negative sepsis and endotoxic shock (Michie, H.R. *et al.*, *Br. J. Surg.* 76:670-671 (1989); Debets, J.M.H. *et al.*, *Second Vienna Shock Forum*, p.463-466 (1989); Simpson, S.Q. *et al.*, *Crit. Care Clin.* 5:27-47 (1989)), including fever, malaise, anorexia, and cachexia. Endotoxin is a potent
10 monocyte/macrophage activator which stimulates production and secretion of TNF (Kornbluth, S.K. *et al.*, *J. Immunol.* 137:2585-2591 (1986)) and other cytokines. Elevated levels of circulating TNF have also been found in patients suffering from Gram-negative sepsis (Waage, A. *et al.*, *Lancet* 1:355-357 (1987); Hammerle, A.F. *et al.*, *Second Vienna Shock Forum* p. 715-718 (1989); Debets,
15 J.M.H. *et al.*, *Crit. Care Med.* 17:489-497 (1989); Calandra, T. *et al.*, *J. Infect. Dis.* 161:982-987 (1990)).

 Neutralizing antisera or mAbs to TNF have been shown in mammals other than man to abrogate adverse physiological changes and prevent death after lethal challenge in experimental endotoxemia and bacteremia. This effect has
20 been demonstrated, e.g., in rodent lethality assays and in primate pathology model systems (Mathison, J.C. *et al.*, *J. Clin. Invest.* 81:1925-1937 (1988); Beutler, B. *et al.*, *Science* 229:869-871 (1985); Tracey, K.J. *et al.*, *Nature* 330:662-664 (1987); Shimamoto, Y. *et al.*, *Immunol. Lett.* 17:311-318 (1988); Silva, A.T. *et al.*, *J. Infect. Dis.* 162:421-427 (1990); Opal, S.M. *et al.*, *J. Infect.*
25 *Dis.* 161:1148-1152 (1990); Hinshaw, L.B. *et al.*, *Circ. Shock* 30:279-292 (1990)). To date, experience with anti-TNF mAb therapy in humans has been limited but shows beneficial therapeutic results, e.g., in arthritis and sepsis. See, e.g., Elliott, M.J. *et al.*, *Baillieres Clin. Rheumatol.* 9:633-52 (1995); Feldmann M, *et al.*, *Ann. N. Y. Acad. Sci. USA* 766:272-8 (1995); van der Poll, T. *et al.*,
30 *Shock* 3:1-12 (1995); Wherry *et al.*, *Crit. Care. Med.* 21:S436-40 (1993); Tracey K.J., *et al.*, *Crit. Care Med.* 21:S415-22 (1993).

-45-

As endokine alpha is believed to exhibit many of the biological effects of TNF, the present invention is further directed antibody-based therapies which involve administering an anti-endokine alpha antibody to a mammalian, preferably human, patient for treating one or more of the above-described disorders. Methods for producing anti-endokine alpha polyclonal and monoclonal antibodies are described in detail above. Such antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding endokine alpha locally or systemically in the body or by direct cytotoxicity of the antibody, *e.g.*, as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended purpose. Amounts and regimens for the administration of antibodies, their fragments or derivatives can be determined readily by those with ordinary skill in the clinical art of treating TNF-related disease.

For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, or buccal routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

Compositions within the scope of this invention include all compositions wherein the antibody, fragment or derivative is contained in an amount effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the

-46-

art. The effective dose is a function of the individual chimeric or monoclonal antibody, the presence and nature of a conjugated therapeutic agent (see below), the patient and his clinical status, and can vary from about 10 µg/kg body weight to about 5000 mg/kg body weight. The preferred dosages comprise 0.1 to 500 mg/kg body wt.

In addition to the pharmacologically active compounds, the new pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Preferably, the preparations, contain from about 0.01 to 99 percent, preferably from about 20 to 75 percent of active compound(s), together with the excipient.

Similarly, preparations of an endokine alpha antibody or fragment of the present invention for parenteral administration, such as in detectably labeled form for imaging or in a free or conjugated form for therapy, include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil such as olive oil, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media, parenteral vehicles including sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, such as those based on Ringer's dextrose, and the like. Preservatives and other additives may also be present, such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like. See, generally, *Remington's Pharmaceutical Science*, 16th ed., Mack Publishing Co., Easton, PA, 1980.

In particular, the antibodies, fragments and derivatives of the present invention are useful for treating a subject having or developing endokine alpha related disorders as described herein. Such treatment comprises parenterally administering a single or multiple doses of the antibody, fragment or derivative, or a conjugate thereof.

-47-

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hemopoietic growth factors, etc., which serve to increase the number or activity of effector cells which interact with the antibodies.

5 Since circulating concentrations of endokine alpha (like TNF) tend to be extremely low, in the range of about 10 pg/ml in non-septic individuals, and reaching about 50 pg/ml in septic patients and above 100 pg/ml in the sepsis syndrome for TNF (Hammerle, A.F. *et al.*, 1989, *supra*) or may be only be detectable at sites of endokine alpha-related disorders, it is preferred to use high
10 affinity and/or potent *in vivo* endokine alpha-inhibiting and/or neutralizing antibodies, fragments or regions thereof, for both endokine alpha immunoassays and therapy of endokine related disorders. Such antibodies, fragments, or regions, will preferably have an affinity for human endokine alpha, expressed as K_a , of at least 10^8 M^{-1} , more preferably, at least 10^9 M^{-1} , such as $5 \times 10^8 \text{ M}^{-1}$, $8 \times 10^8 \text{ M}^{-1}$, $2 \times 10^9 \text{ M}^{-1}$, $4 \times 10^9 \text{ M}^{-1}$, $6 \times 10^9 \text{ M}^{-1}$, $8 \times 10^9 \text{ M}^{-1}$.
15

Preferred for human therapeutic use are high affinity murine and murine/human or human/human chimeric antibodies, and fragments, regions and derivatives having potent *in vivo* endokine-inhibiting and/or neutralizing activity, according to the present invention, *e.g.*, that block endokine-induced IL-1, IL-6
20 or TNF secretion, procoagulant activity, expression of cell adhesion molecules such as ELAM-1 and ICAM-1 and mitogenic activity, *in vivo*, *in situ*, and *in vitro*.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way
25 of illustration and are not intended as limiting.

Examples

Example 1: Expression and Purification of Mature Endokine Alpha in E. coli

5 The DNA sequence encoding the mature endokine alpha protein in the deposited cDNA clone is amplified using PCR oligonucleotide primers specific to the amino terminal sequences of the endokine alpha protein. Additional nucleotides containing restriction sites to facilitate cloning were added to the 5' and 3' sequences, respectively.

10 The 5' oligonucleotide primer had the sequence GCG CCA TGG CTA AGT TTG GAC CAT (SEQ ID NO:5) containing the underlined *Nco* I restriction site.

The 3' primer had the sequence GCG AAG CTT TCA AGT CTC TAG GAG ATG (SEQ ID NO:6) containing the underlined *Hind*III restriction site.

15 The restriction sites are convenient to restriction enzyme sites in the bacterial expression vector pQE60, which is used for bacterial expression in M15/rep4 host cells in these examples. (Qiagen, Inc., Chatsworth, CA, 91311). pQE60 encodes ampicillin antibiotic resistance ("Amp") and contains a bacterial origin of replication ("ori"), an IPTG inducible promoter, a ribosome binding site ("RBS"), a 6-His tag and restriction enzyme sites.

20 The amplified endokine alpha protein DNA and the vector pQE60 both are digested with *Nco*I and *Hind*III and the digested DNAs are then ligated together. Insertion of the endokine alpha protein DNA into the restricted pQE60 vector places the endokine alpha protein coding region downstream of and operably linked to the vector's IPTG-inducible promoter and in-frame with an initiating ATG appropriately positioned for translation of endokine alpha protein.

25 The ligation mixture is transformed into competent *E. coli* cells using standard procedures. Such procedures are described in Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). *E. coli* strain M15/rep4,

-49-

containing multiple copies of the plasmid pREP4, which expresses lac repressor and confers kanamycin resistance ("Kan"), is used in carrying out the illustrative example described here. This strain, which is only one of many that are suitable for expressing endokine alpha protein, is available commercially from Qiagen.

5 Transformants are identified by their ability to grow on LB plates in the presence of ampicillin and kanamycin. Plasmid DNA is isolated from resistant colonies and the identity of the cloned DNA confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight ("O/N") in liquid culture in LB media supplemented with both ampicillin (100 µg/ml) and
10 kanamycin (25 µg/ml).

The O/N culture is used to inoculate a large culture, at a dilution of approximately 1:100 to 1:250. The cells are grown to an optical density at 600 NM ("OD600") of between 0.4 and 0.6. Isopropyl-B-D-thiogalactopyranoside ("IPTG") is then added to a final concentration of 1 mM
15 to induce transcription from *lac* repressor sensitive promoters, by inactivating the *lacI* repressor. Cells subsequently are incubated further for 3 to 4 hours. Cells then are harvested by centrifugation and disrupted, by standard methods. Inclusion bodies are purified from the disrupted cells using routine collection techniques, and protein is solubilized from the inclusion bodies into 8M urea.
20 The 8M urea solution containing the solubilized protein is passed over a PD-10 column in 2X phosphate-buffered saline ("PBS"), thereby removing the urea, exchanging the buffer and refolding the protein. The protein is purified by a further step of chromatography to remove endotoxin. Then, it is sterile filtered. The sterile filtered protein preparation is stored in 2X PBS.

25 ***Example 2: Cloning and Expression of Mature Endokine Alpha in a Baculovirus Expression System***

The cDNA sequence encoding the entire endokine alpha protein in the deposited clone is amplified using PCR oligonucleotide primers corresponding
30 to 5' and 3' regions of the gene.

-50-

The 5' primer has the sequence GC GGA TCC CGA GAC TGC TAA GGA GCC (SEQ ID NO:7) containing the underlined *Bam*HI restriction enzyme site and containing a sequence complementary to that encoding a portion of the endokine alpha protein in FIG. 1.

5 The 3' primer has the sequence GC GGA TCC CTA GGA GAT GAA TTG GGG ATT TG (SEQ ID NO:8) containing the underlined *Bam*HI restriction site and containing a sequence complementary to that encoding a portion of the endokine alpha protein in FIG. 1.

10 The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("Geneclean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with *Bam*HI and again is purified on a 1% agarose gel. This fragment is designated herein F2.

15 The vector pA2-GP is used to express the endokine alpha protein in the baculovirus expression system, using standard methods, as described in Summers *et al.*, *A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures*, Texas Agricultural Experimental Station Bulletin No. 1555 (1987). This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites. The signal peptide of AcMNPV gp67, including the
20 N-terminal methionine, is located just upstream of a *Bam*HI site. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For an easy selection of recombinant virus, the beta-galactosidase gene from *E. coli* is inserted in the same orientation as the polyhedrin promoter and is followed by the polyadenylation signal of the
25 polyhedrin gene. The polyhedrin sequences are flanked at both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate viable virus that express the cloned polynucleotide.

30 Many other baculovirus vectors could be used in place of pA2-GP, such as pAc373, pVL941 and pAcIM1 provided, as those of skill readily will appreciate, that construction provides appropriately located signals for transcription, translation, trafficking and the like, such as an in-frame AUG and

-51-

a signal peptide, as required. Such vectors are described in Luckow *et al.*, *Virology* 170: 31-39, among others.

The plasmid is digested with the restriction enzyme XbaI and then is dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("Geneclean" BIO 101 Inc., La Jolla, Ca.). This vector DNA is designated herein "V2".

Fragment F2 and the dephosphorylated plasmid V2 are ligated together with T4 DNA ligase. *E. coli* HB101 cells are transformed with ligation mix and spread on culture plates. Bacteria are identified that contain the plasmid with the human endokine alpha gene by digesting DNA from individual colonies using XbaI and then analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

5 µg of the plasmid is co-transfected with 1.0 µg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA.), using the lipofection method described by Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84: 7413-7417 (1987). 1 µg of BaculoGold™ virus DNA and 5 µg of the plasmid are mixed in a sterile well of a microtiter plate containing 50 µl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards 10 µl Lipofectin plus 90 µl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is rocked back and forth to mix the newly added solution. The plate is then incubated for 5 hours at 27°C. After 5 hours the transfection solution is removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. The plate is put back into an incubator and cultivation is continued at 27°C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, cited above. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy

-52-

identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10).

5 Four days after serial dilution, the virus is added to the cells. After appropriate incubation, blue stained plaques are picked with the tip of an Eppendorf pipette. The agar containing the recombinant viruses is then resuspended in an Eppendorf tube containing 200 μ l of Grace's medium. The agar is removed by a brief centrifugation and the supernatant containing the
10 recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4°C. A clone containing properly inserted hESSB I, II and III is identified by DNA analysis including restriction mapping and sequencing of this plasmid.

15 Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus at a multiplicity of infection ("MOI") of about 2 (about 1 to about 3). Six hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Gaithersburg). 42 hours
20 later, 5 μ Ci of 35 S-methionine and 5 μ Ci 35 S-cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then they are harvested by centrifugation, lysed and the labeled proteins are visualized by SDS-PAGE and autoradiography.

Example 3: Cloning and Expression in CHO Cells

25 The vector pC1 is used for the expression of endokine alpha protein. Plasmid pC1 is a derivative of the plasmid pSV2-dhfr [ATCC Accession No. 37146]. Both plasmids contain the mouse DHFR gene under control of the SV40 early promoter. Chinese hamster ovary- or other cells lacking dihydrofolate activity that are transfected with these plasmids can be selected by growing the

-53-

cells in a selective medium (alpha minus MEM, Life Technologies) supplemented with the chemotherapeutic agent methotrexate. The amplification of the DHFR genes in cells resistant to methotrexate (MTX) has been well documented (*see, e.g., Alt, F.W. et al. J. Biol. Chem. 253:1357-1370 (1978), Hamlin, J.L. and Ma, C., Biochim. et Biophys. Acta, 1097:107-143 (1990), Page, M.J. and Sydenham, M.A., Biotechnology 9:64-68 (1991)*). Cells grown in increasing concentrations of MTX develop resistance to the drug by overproducing the target enzyme, DHFR, as a result of amplification of the DHFR gene. If a second gene is linked to the DHFR gene it is usually co-amplified and over-expressed. It is state of the art to develop cell lines carrying more than 1,000 copies of the genes. Subsequently, when the methotrexate is withdrawn, cell lines contain the amplified gene integrated into the chromosome(s).

Plasmid pC1 contains for the expression of the gene of interest a strong promoter of the long terminal repeat (LTR) of the Rouse Sarcoma Virus (Cullen, *et al., Molecular and Cellular Biology, 438-4470 (March 1985)*) plus a fragment isolated from the enhancer of the immediate early gene of human cytomegalovirus (CMV) (Boshart *et al., Cell 41:521-530 (1985)*). Downstream of the promoter are the following single restriction enzyme cleavage sites that allow the integration of the genes: BamHI, PvuII, and NruI. Behind these cloning sites the plasmid contains translational stop codons in all three reading frames followed by the 3' intron and the polyadenylation site of the rat preproinsulin gene. Other high efficient promoters can also be used for the expression, e.g., the human β -actin promoter, the SV40 early or late promoters or the long terminal repeats from other retroviruses, e.g., HIV and HTLV. For the polyadenylation of the mRNA other signals, e.g., from the human growth hormone or globin genes can be used as well.

Stable cell lines carrying a gene of interest integrated into the chromosomes can also be selected upon co-transfection with a selectable marker such as gpt, G418 or hygromycin. It is advantageous to use more than one selectable marker in the beginning, e.g. G418 plus methotrexate.

-54-

The plasmid pC1 is digested with the restriction enzyme *Bam*HI and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

5 The DNA sequence encoding endokine alpha, ATCC No. 97640, is amplified using PCR oligonucleotide primers corresponding to the 5' and 3' sequences of the gene:

10 The 5' primer has the sequence 5' GCG GGA TCC GCC ATC ATG CCT TTA AGC CAT TC 3' (SEQ ID NO:9) containing the underlined *Bam*HI restriction enzyme site followed by bases of the sequence of endokine alpha of FIG. 1 (SEQ ID NO:1). Inserted into an expression vector, as described below, the 5' end of the amplified fragment encoding human endokine alpha provides an efficient signal peptide. An efficient signal for initiation of translation in eukaryotic cells, as described by Kozak, M., *J. Mol. Biol.* 196:947-950 (1987) is appropriately located in the vector portion of the construct.

15 The 3' primer has the sequence 5' GC GGA TCC CTA GGA GAT GAA TTG GGG ATT TG 3' (SEQ ID NO:10) containing the *Asp*718 restriction followed by nucleotides complementary to the endokine alpha coding sequence set out in FIG. 1 (SEQ ID NO:1), including the stop codon.

20 The amplified fragments are isolated from a 1% agarose gel as described above and then digested with the endonucleases *Bam*HI and *Asp*718 and then purified again on a 1% agarose gel.

25 The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E.coli* HB101 cells are then transformed and bacteria identified that contained the plasmid pC1 inserted in the correct orientation using the restriction enzyme *Bam*HI. The sequence of the inserted gene is confirmed by DNA sequencing.

Transfection of CHO-DHFR-cells

Chinese hamster ovary cells lacking an active DHFR enzyme are used for transfection. 5 µg of the expression plasmid C1 are cotransfected with 0.5 µg of

-55-

the plasmid pSVneo using the lipofecting method (Felgner *et al.*, *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the gene neo from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) and cultivated from 10-14 days. After this period, single clones are trypsinized and then seeded in 6-well petri dishes using different concentrations of methotrexate (25 nM, 50 nM, 100 nM, 200 nM, 400 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (500 nM, 1 μ M, 2 μ M, 5 μ M). The same procedure is repeated until clones grow at a concentration of 100 μ M.

The expression of the desired gene product is analyzed by Western blot analysis and SDS-PAGE.

Example 4: Tissue distribution of Endokine Alpha expression

Northern blot analysis was carried out to examine the levels of expression of the gene encoding the endokine alpha protein in human tissues, using methods described by, among others, Sambrook *et al.*, *supra*. A cDNA probe containing the entire nucleotide sequence of the endokine alpha protein of the present invention (SEQ ID NO:1) was labeled with 32 P using the *rediprime*TM DNA labeling system (Amersham Life Science), according to manufacturer's instructions. After labelling, the probe was purified using a CHROMA SPIN-100TM column (Clontech Laboratories, Inc.), according to manufacturer's protocol number PT1200-1. The purified labelled probe was then used to examine various human tissues for the expression of the gene encoding the endokine alpha protein.

Multiple Tissue Northern (MTN) blots containing various human tissues (H) or human immune system tissues (IM) were obtained from Clontech and were examined with labelled probe using *ExpressHyb*TM Hybridization Solution (Clontech) according to manufacturer's protocol number PT1190-1. Following

-56-

hybridization and washing, the blots were mounted and exposed to film at -70°C overnight, and films developed according to standard procedures.

Expression of the gene encoding an endokine alpha protein of the present invention was detected in human brain striatum and pancreas tissue.

5 It will be clear that the invention may be practiced otherwise than as particularly described in the foregoing description and examples.

Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

10 The disclosures of all patents, patent applications, and publications referred to herein are hereby incorporated by reference.

-57-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Human Genome Sciences, Inc.
9410 Key West Avenue
Rockville, MD 20850
United States of America
APPLICANTS/INVENTORS: Yu, Guo-Liang
Ni, Jian
Rosen, Craig A.

(ii) TITLE OF INVENTION: Human Endokine Alpha

(iii) NUMBER OF SEQUENCES: 10

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: STERNE, KESSLER, GOLDSTEIN & FOX, P.L.L.C
(B) STREET: 1100 NEW YORK AVE., NW, SUITE 600
(C) CITY: WASHINGTON
(D) STATE: DC
(E) COUNTRY: USA
(F) ZIP: 20005-3934

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version #1.30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: PCT To be assigned
(B) FILING DATE: Herewith
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Goldstein, Jorge A.
(B) REGISTRATION NUMBER: 29,021
(C) REFERENCE/DOCKET NUMBER: 1488.047PC00

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 202-371-2600
(B) TELEFAX: 202-371-2540

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1809 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: both
(D) TOPOLOGY: both

(ii) MOLECULE TYPE: cDNA

-58-

(ix) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 53..559

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GTTTTCCACA GCTCTCATTT CTCCAAAAAT GTGTTTGAGC CACTTGGAAA AT ATG	55
Met	
1	
CCT TTA AGC CAT TCA AGA ACT CAA GGA GCT CAG AGA TCA TCC TGG AAG	103
Pro Leu Ser His Ser Arg Thr Gln Gly Ala Gln Arg Ser Ser Trp Lys	
5 10 15	
CTG TGG CTC TTT TGC TCA ATA GTT ATG TTG CTA TTT CTT TGC TCC TTC	151
Leu Trp Leu Phe Cys Ser Ile Val Met Leu Leu Phe Leu Cys Ser Phe	
20 25 30	
AGT TGG CTA ATC TTT ATT TTT CTC CAA TTA GAG ACT GCT AAG GAG CCC	199
Ser Trp Leu Ile Phe Ile Phe Leu Gln Leu Glu Thr Ala Lys Glu Pro	
35 40 45	
TGT ATG GCT AAG TTT GGA CCA TTA CCC TCA AAA TGG CAA ATG GCA TCT	247
Cys Met Ala Lys Phe Gly Pro Leu Pro Ser Lys Trp Gln Met Ala Ser	
50 55 60 65	
TCT GAA CCT CCT TGC GTG AAT AAG GTG TCT GAC TGG AAG CTG GAG ATA	295
Ser Glu Pro Pro Cys Val Asn Lys Val Ser Asp Trp Lys Leu Glu Ile	
70 75 80	
CTT CAG AAT GGC TTA TAT TTA ATT TAT GGC CAA GTG GCT CCC AAT GCA	343
Leu Gln Asn Gly Leu Tyr Leu Ile Tyr Gly Gln Val Ala Pro Asn Ala	
85 90 95	
AAC TAC AAT GAT GTA GCT CCT TTT GAG GTG CGG CTG TAT AAA AAC AAA	391
Asn Tyr Asn Asp Val Ala Pro Phe Glu Val Arg Leu Tyr Lys Asn Lys	
100 105 110	
GAC ATG ATA CAA ACT CTA ACA AAC AAA TCT AAA ATC CAA AAT GTA GGA	439
Asp Met Ile Gln Thr Leu Thr Asn Lys Ser Lys Ile Gln Asn Val Gly	
115 120 125	
GGG ACT TAT GAA TTG CAT GTT GGG GAC ACC ATA GAC TTG ATA TTC AAC	487
Gly Thr Tyr Glu Leu His Val Gly Asp Thr Ile Asp Leu Ile Phe Asn	
130 135 140 145	
TCT GAG CAT CAG GTT CTA AAA AAT AAT ACC TAC TGG GGT ATC ATT TTA	535
Ser Glu His Gln Val Leu Lys Asn Asn Thr Tyr Trp Gly Ile Ile Leu	
150 155 160	
CTG GCA AAT CCC CAA TTC ATC TCC TAGAGACTTG ATTTGATCTC CTCATTCCCT	589
Leu Ala Asn Pro Gln Phe Ile Ser	
165	

-59-

TCAGCACATG TAGAGGTGCC AGTGGGTGGA TTGGAGGGAG AAGATATTCA ATTTCTAGAG	649
TTTCTGTCTA CAAAAATCAA CACAAACAGA ACTCCTCTGC ACGTGAATTT TCATCTATCA	709
TGCATCTGAA AGAGACTCAG GGGAAAAGCC AAAGACTTTT GGTTGGATCT GCAGAGATAC	769
TTCTAATCCA TGATAAAACA AATATGGATG ACAGAGGACA TGTGCTTTTC AAAGAATCTT	829
TATAATTCTT GAATTCATGA GTGGAAAAAT GGAGTTCTAT TCCCATGGAA GATTTACCTG	889
GTACAAAAAG GATCTGGGGC AGTAGCCTGG CTTTGTTCTC ATATTCTTGG GCTGCTGTAA	949
TTCTCTTCTC ATACTCCCAT CTTCTGAGAC CCTCCCAATA AAAAGTAGAC TGATAGGATG	1009
GCCAGATATG CCTACCATAC CCTACTTTAG ATATGGTGGT GTTAGAAGAT AAAGAACAAT	1069
CTGAACTATT GGAATAGAGG TACAAGTGGC ATAAAATGGA ATGTACGCTA TCTGGAAATT	1129
TCTTGGTTTT ATCTTCCTCA GGATGCAGGG TGCTTTAAAA AGCCTTATCA AAGGAGTCAT	1189
TCCACCCTCA CGTAGAGCTT TGTGAGAACT TACTGTTGGT GTGTGTGTCT AAACATTGCT	1249
AATTAAAGAA AGAGTAACCA TTAGTAATCA TTAGGTTTAA CCCCAGAATG GTATTATCAT	1309
TACGATTATG TCATGTAATG ATTTAGTATT TTTAGCTAGC TTTCCACAGT TTGCAAAGTG	1369
CTTGTAAAAC AGTTAGCAAT TCTATGAAGT TAATTGGGCA GGCATTTGGG GGAAAATTTT	1429
AGTTGAGAAT GTGATAGCAT AGCATAGCCA ACTTTCCTCA ACTCATAGGA CAAGTGACTA	1489
CAAGGCAATG GGTAGTCCCC TGCATTGCAC TGTCTCAGCT TTAGAATTGT TATTTCTGCT	1549
ATCGTTATAA GACTCTAAAA CTTAGCGAAT TCACTTTTCA GGAAGCATAT TCCCCTTTAG	1609
CCCGGTGAGC AGAGTGAAGC TACAACAGAT CTTTCCTTTA CCAGCACACT TTTTTTTTTT	1669
TCCCCTGAAT CAGGGAGATC CAGGATGCTG TTCAGGCCTT ATCCCAACCA AATCCCCCTC	1729
TTCTTTGCAG GGCCCATCTT AGTCAAATGT GCTAACTTCT AAAATAATAA ATAGCACTAA	1789
TTCAAAAAAA AAAAAAAAAA	1809

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 169 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Pro Leu Ser His Ser Arg Thr Gln Gly Ala Gln Arg Ser Ser Trp

SUBSTITUTE SHEET (RULE 26)

-60-

1	5	10	15												
Lys	Leu	Trp	Leu	Phe	Cys	Ser	Ile	Val	Met	Leu	Leu	Phe	Leu	Cys	Ser
	20							25						30	
Phe	Ser	Trp	Leu	Ile	Phe	Ile	Phe	Leu	Gln	Leu	Glu	Thr	Ala	Lys	Glu
	35						40					45			
Pro	Cys	Met	Ala	Lys	Phe	Gly	Pro	Leu	Pro	Ser	Lys	Trp	Gln	Met	Ala
	50					55					60				
Ser	Ser	Glu	Pro	Pro	Cys	Val	Asn	Lys	Val	Ser	Asp	Trp	Lys	Leu	Glu
	65				70					75					80
Ile	Leu	Gln	Asn	Gly	Leu	Tyr	Leu	Ile	Tyr	Gly	Gln	Val	Ala	Pro	Asn
			85						90					95	
Ala	Asn	Tyr	Asn	Asp	Val	Ala	Pro	Phe	Glu	Val	Arg	Leu	Tyr	Lys	Asn
			100					105						110	
Lys	Asp	Met	Ile	Gln	Thr	Leu	Thr	Asn	Lys	Ser	Lys	Ile	Gln	Asn	Val
	115						120					125			
Gly	Gly	Thr	Tyr	Glu	Leu	His	Val	Gly	Asp	Thr	Ile	Asp	Leu	Ile	Phe
	130					135					140				
Asn	Ser	Glu	His	Gln	Val	Leu	Lys	Asn	Asn	Thr	Tyr	Trp	Gly	Ile	Ile
	145				150					155					160
Leu	Leu	Ala	Asn	Pro	Gln	Phe	Ile	Ser							
					165										

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 233 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: not relevant
 - (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met	Ser	Thr	Glu	Ser	Met	Ile	Arg	Asp	Val	Glu	Leu	Ala	Glu	Glu	Ala
1				5					10					15	
Leu	Pro	Lys	Lys	Thr	Gly	Gly	Pro	Gln	Gly	Ser	Arg	Arg	Cys	Leu	Phe
		20						25					30		
Leu	Ser	Leu	Phe	Ser	Phe	Leu	Ile	Val	Ala	Gly	Ala	Thr	Thr	Leu	Phe
	35						40					45			

-61-

Cys Leu Leu His Phe Gly Val Ile Gly Pro Gln Arg Glu Glu Ser Pro
 50 55 60
 Arg Asp Leu Ser Leu Ile Ser Pro Leu Ala Gln Ala Val Arg Ser Ser
 65 70 75 80
 Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val Val Ala Asn Pro
 85 90 95
 Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg Ala Asn Ala Leu
 100 105 110
 Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu Val Val Pro Ser
 115 120 125
 Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe Lys Gly Gln Gly
 130 135 140
 Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile Ser Arg Ile Ala
 145 150 155 160
 Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala Ile Lys Ser Pro
 165 170 175
 Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys Pro Trp Tyr Glu
 180 185 190
 Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys Gly Asp Arg Leu
 195 200 205
 Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp Phe Ala Glu Ser Gly
 210 215 220
 Gln Val Tyr Phe Gly Ile Ile Ala Leu
 225 230

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 205 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: not relevant
- (D) TOPOLOGY: not relevant

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Thr Pro Pro Glu Arg Leu Phe Leu Pro Arg Val Cys Gly Thr Thr
 1 5 10 15
 Leu His Leu Leu Leu Leu Gly Leu Leu Leu Val Leu Leu Pro Gly Ala

-62-

20					25					30					
Gln	Gly	Leu	Pro	Gly	Val	Gly	Leu	Thr	Pro	Ser	Ala	Ala	Gln	Thr	Ala
		35					40					45			
Arg	Gln	His	Pro	Lys	Met	His	Leu	Ala	His	Ser	Thr	Leu	Lys	Pro	Ala
		50					55					60			
Ala	His	Leu	Ile	Gly	Asp	Pro	Ser	Lys	Gln	Asn	Ser	Leu	Leu	Trp	Arg
							70					75			80
Ala	Asn	Thr	Asp	Arg	Ala	Phe	Leu	Gln	Asp	Gly	Phe	Ser	Leu	Ser	Asn
				85					90					95	
Asn	Ser	Leu	Leu	Val	Pro	Thr	Ser	Gly	Ile	Tyr	Phe	Val	Tyr	Ser	Gln
			100					105					110		
Val	Val	Phe	Ser	Gly	Lys	Ala	Tyr	Ser	Pro	Lys	Ala	Pro	Ser	Ser	Pro
			115					120				125			
Leu	Tyr	Leu	Ala	His	Glu	Val	Gln	Leu	Phe	Ser	Ser	Gln	Tyr	Pro	Phe
		130					135					140			
His	Val	Pro	Leu	Leu	Ser	Ser	Gln	Lys	Met	Val	Tyr	Pro	Gly	Leu	Gln
		145					150					155			160
Glu	Pro	Trp	Leu	His	Ser	Met	Tyr	His	Gly	Ala	Ala	Phe	Gln	Leu	Thr
			165						170					175	
Gln	Gly	Asp	Gln	Leu	Ser	Thr	His	Thr	Asp	Gly	Ile	Pro	His	Leu	Val
			180						185				190		
Leu	Ser	Pro	Ser	Thr	Val	Phe	Phe	Gly	Ala	Phe	Ala	Leu			
		195					200					205			

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 24 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GCGCCATGGC TAAGTTTGGA CCAT

24

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:

-63-

- (A) LENGTH: 27 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GCGAAGCTTT CAAGTCTCTA GGAGATG

27

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GCGGATCCCG AGACTGCTAA GGAGCC

26

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GCGGATCCCT AGGAGATGAA TTGGGGATTT G

31

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

-64-

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GCGGGATCCG CCATCATGCC TTAAAGCCAT TC

32

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 31 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCGGATCCCT AGGAGATGAA TTGGGGATTT G


31

-64.1-

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>5</u> , line <u>20</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution AMERICAN TYPE CULTURE COLLECTION	
Address of depositary institution (including postal code and country) 12301 Parklawn Drive Rockville, Maryland 20852 United States of America	
Date of deposit June 27, 1996	Accession Number ATCC 97640
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
DNA PLASMID HG944451	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input checked="" type="checkbox"/> This sheet was received with the international application
Authorized officer 

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

-65-

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:

5 (a) a nucleotide sequence encoding the full-length endokine alpha polypeptide having the complete amino acid sequence in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640;

(b) a nucleotide sequence encoding the extracellular endokine alpha polypeptide having an amino acid sequence from about residue 44 to about residue 169 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640;

10 (c) a nucleotide sequence encoding the endokine alpha transmembrane domain having an amino acid sequence from about residue 18 to about residue 43 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640;

15 (d) a nucleotide sequence encoding the endokine alpha intracellular domain having an amino acid sequence from about residue 1 to about residue 17 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640; and

20 (e) a nucleotide sequence complementary to any of the nucleotide sequences in (a), (b), (c) or (d).

2. An isolated nucleic acid molecule comprising a polynucleotide which hybridizes under stringent hybridization conditions to a polynucleotide having a nucleotide sequence identical to a nucleotide sequence in (a), (b), (c), (d) or (e) of claim 1 wherein said polynucleotide which does not hybridize under stringent hybridization conditions to a polynucleotide having a nucleotide sequence consisting of only A residues or of only T residues.

-66-

3. An isolated nucleic acid molecule comprising a polynucleotide which encodes the amino acid sequence of an epitope-bearing portion of an endokine alpha polypeptide having an amino acid sequence in (a), (b), (c), (d), or (e) of claim 1.

5 4. The isolated nucleic acid molecule of claim 3, which encodes an epitope-bearing portion of an endokine alpha polypeptide selected from the group consisting of: a polypeptide comprising amino acid residues from about 44 to about 158 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 44 to about 54 in FIG. 1 (SEQ ID NO:2); a polypeptide
10 comprising amino acid residues from about 57 to about 68 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 69 to about 78 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 94 to about 105 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 108 to about 132 in FIG. 1 (SEQ ID NO:2); and a
15 polypeptide comprising amino acid residues from about 148 to about 158 in FIG. 1 (SEQ ID NO:2).

5. The isolated nucleic acid molecule of claim 1, wherein said polynucleotide is DNA.

20 6. The isolated nucleic acid molecule of claim 1, wherein said polynucleotide is RNA.

7. A method for making a recombinant vector comprising inserting an isolated nucleic acid molecule of claim 1 into a vector.

8. A recombinant vector produced by the method of claim 7.

25 9. A method of making a recombinant host cell comprising introducing the recombinant vector of claim 8 into a host cell.

-67-

10 A recombinant host cell produced by the method of claim 9.

5 11. A recombinant method for producing an endokine alpha polypeptide, comprising culturing the recombinant host cell of claim 10 under conditions such that said polypeptide is expressed and recovering said polypeptide.

12. An isolated polypeptide having an amino acid sequence at least 95% identical to a sequence selected from the group consisting of:

10 (a) the amino acid sequence of the full-length endokine alpha polypeptide in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640;

(b) the amino acid sequence of the extracellular endokine alpha polypeptide from about residue 44 to about residue 169 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640;

15 (c) the amino acid sequence of the endokine alpha transmembrane domain from about residue 18 to about residue 43 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640;

20 (d) the amino acid sequence of the endokine alpha intracellular domain from about residue 1 to about residue 17 in FIG. 1 (SEQ ID NO:2) or as encoded by the cDNA clone contained in ATCC Deposit No. 97640; and

(e) the amino acid sequence of an epitope-bearing portion of any one of the polypeptides of (a), (b), (c), or (d).

25 13. An isolated polypeptide comprising an epitope-bearing portion of endokine alpha, wherein said portion is selected from the group consisting of:
a polypeptide comprising amino acid residues from about 44 to about 158 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 44 to about 54 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 57 to about 68 in FIG. 1 (SEQ ID NO:2); a polypeptide

-68-

comprising amino acid residues from about 69 to about 78 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 94 to about 105 in FIG. 1 (SEQ ID NO:2); a polypeptide comprising amino acid residues from about 108 to about 132 in FIG. 1 (SEQ ID NO:2); and a polypeptide comprising amino acid residues from about 148 to about 158 in FIG. 1 (SEQ ID NO:2).

14. An isolated antibody or antibody fragment that binds specifically to an endokine alpha polypeptide of claim 12.

15. A method for treating an individual in need of a decreased level of endokine alpha activity, comprising
administering to said individual a composition comprising the isolated antibody or antibody fragment of claim 14.

16. A diagnostic method, comprising:
(a) assaying endokine alpha gene expression level in mammalian cells or body fluid; and
(b) comparing said endokine alpha gene expression level with a standard endokine alpha gene expression level, whereby an increase or decrease in said endokine alpha gene expression level compared to said standard is indicative of a TNF-related disorder.

1/4

1 GTTTTCCACAGCTCTCATTTCTCCAAAAATGTGTTTGAGCCACTTGAAAAAT
ATGCCTTTAAGCCATTCAAGAACTCAAGGAGCTCAGAGATCATCCTGGAAGCTGTGGCTC
M P L S H S R T Q G A Q R S S W K L W L

61 TTTTGCTCAATAGTTATGTTGCTATTTCTTTGCTCCTTCAGTTGGCTAATCTTTATTTT
F C S I V M L L F L C S F S W L I F I F

121 CTCCAATTAGAGACTGCTAAGGAGCCCTGTATGGCTAAGTTTGGACCATTACCCTCAAAA
L Q L E T A K E P C M A K F G P L P S K

181 TGGCAAATGGCATCTTCTGAACCTCCTTGCGTGAATAAGGTGTCTGACTGGAAGCTGGAG
W Q M A S S E P P C V N K V S D W K L E

241 ATACTTCAGAATGGCTTATATTTAATTTATGGCCAAGTGGCTCCCAATGCAAACCTACAAT
I L Q N G L Y L I Y G Q V A P N A N Y N

301 GATGTAGCTCCTTTTGAGGTGCGGCTGTATAAAAAACAAAGACATGATACAACTCTAACA
D V A P F E V R L Y K N K D M I Q T L T

361 AACAAATCTAAATCCAAATGTAGGAGGGACTTATGAATTGCATGTTGGGGACACCATA
N K S K I Q N V G G T Y E L H V G D T I

421 GACTTGATATTCAACTCTGAGCATCAGGTTCTAAAAAATAATACCTACTGGGGTATCATT
D L I F N S E H Q V L K N N T Y W G I I

481 TTA CTGGCAAATCCCAATTCATCTCCTAGAGACTTGATTTGATCTCCTCATTCCCTTCA
L L A N P Q F I S

GCACATGTAGAGGTGCCAGTGGGTGGATTGGAGGGAGAAGATATTCAATTTCTAGAGTTT
CTGTCTACAAAAATCAACACAAACAGAACTCCTCTGCACGTGAATTTTCATCTATCATGC
ATCTGAAAGAGACTCAGGGGAAAAGCCAAAGACTTTTGGTTGGATCTGCAGAGATACTTC
TAATCCATGATAAAACAAATATGGATGACAGAGGACATGTGCTTTTCAAAGAATCTTTAT
AATTCTTGAATTCATGAGTGGAAAAATGGAGTTCTATTCCCATGGAAGATTTACCTGGTA
CAAAAAGGATCTGGGGCAGTAGCCTGGCTTTGTTCTCATATTCTTGGGCTGCTGTAATTC
TCTTCTCATACTCCCATCTTCTGAGACCCTCCCAATAAAAAGTAGACTGATAGGATGGCC
AGATATGCCTACCATACCCTACTTTAGATATGGTGGTGTTAGAAGATAAAGAACAATCTG
AACTATTGGAATAGAGGTACAAGTGGCATAAAATGGAATGTACGCTATCTGGAAATTTCT
TGGTTTTATCTTCCTCAGGATGCAGGGTGCTTTAAAAAGCCTTATCAAAGGAGTCATTCC
ACCCTCACGTAGAGCTTTGTGAGAACTTACTGTTGGTGTGTGTGTCTAAACATTGCTAAT
TAAAGAAAGAGTAACCATTAGTAATCATTAGGTTTAAACCCAGAATGGTATTATCATTAC
GATTATGTCATGTAATGATTTAGTATTTTAGCTAGCTTTCCACAGTTTGCAAAGTGCTT

2/4

GTAAACAGTTAGCAATTCTATGAAGTTAATTGGGCAGGCATTTGGGGGAAAATTTTAGT
TGAGAATGTGATAGCATAGCATAGCCTTCTCACTCATAGGACAAGTGACTACAA
GGCAATGGGTAGTCCCCTGCATTGCACTGTCTCAGCTTTAGAATTGTTATTTCTGCTATC
GTTATAAGACTCTAAACTTAGCGAATTCACCTTTTCAGGAAGCATATCCCCTTTAGCCC
GGTGAGCAGAGTGAAGCTACAACAGATCTTTCCTTTACCAGCACACTTTTTTTTTTTTCC
CCTGAATCAGGGAGATCCAGGATGCTGTTTCAGGCCTTATCCCAACCAAATTCCCCTCTTC
TTTGCAGGGCCCATCTTAGTCAAATGTGCTAACTTCTAAAATAATAAATAGCACTAATTC
AAAAAAAAAAAAAAAAAAAA

FIG.1 B

3/4

```

1 MS-TESMIRDVELAEELPKKTG-----GPGGSRRCFL-----SLFSELIIV--A TNFalpha
1 MT-PPERL-----FLPRVCG-----TT-----HL-----LLGLLVLLP TNFbeta
1 MPLSHSRT-----QGAQRSSWKLWLFCSIVML-----FLCSFSWLTETELDL--E Endokine alpha

43 GATTLCCLHFGVIGPQREESPRDLSLISPLAQAVRSSRT-----PSDKPMHVVANPQA TNFalpha
31 GAQGL-----PGVGLTPSAAGTARQHPKHLAHSTLKPAHLIGDPSK TNFbeta
45 TAKE-----PCMAKFGPLP-----SKVQM-----ASSEPCV Endokine alpha

99 EGGLQMLNRRANALLANGVELRDNLVMPSEGLYLYSQVLFKGG-----CPSTHVLITHT TNFalpha
74 QNSLLVRANTDRAFLDDGSLSNNSLLVPTSGIYFVYSQVVEGKAYSPKAPSSPLYLAH TNFbeta
72 NKVSDMKLEILQ-----NGLYLYGQV-----APNAN-----Endokine alpha

155 TISRIAVSYDTKVNLLSAIKSPCQRETPEGAEAKPWYEPITYLGMFQLEKGRLSAEINR TNFalpha
134 EVQLFSSQYPFHVPILLSQK-----MVYPGLQEPWLSMYHGAFLQTGGDLSTHTDG TNFbeta
99 ----YNDVAPFEVRLYKNKD-----MIQTLTNKSKIQNV--GGTYELHVGDTIDLIFNS Endokine alpha

215 PDYIDFAESGQVYFGTIAL-----TNFalpha
188 IPHLVLS-PSIVFGAFAL-----TNFbeta
147 -EHQVLK-NNT-YAGIILLANPQFIS-----Endokine alpha

```

FIG.2




FIG.3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13282**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C12P 21/06; C12N 1/20, 15/00; C07H 21/04; C07K 1/00, 14/52

US CL : 435/69.5, 69.1, 252.3, 320.1; 536/23.5; 530/351, 350

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/69.5, 69.1, 252.3, 320.1; 536/23.5; 530/351, 350

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Sequence search of Seq. ID No 1 and it's complement, and ID No 2 . Database search for "endokine"

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 5,288,852 A (Yamada et al.) 22 February 1994, see column 4, lines 11-65 and sequence comparison.	1-13 ----- 1-13
A	Database EST-STN on MASPARG search, WashU-Merck EST Project, (St. Louis, Mo, USA) , No. R38487, HILLIER et al., "yf60c04.s1 Homo sapiens cDNA clone 26749 3', 04 May 1995, see sequence alignment.	1-13
A	Database EST-STN on MASPARG search, WashU-Merck EST Project, (St. Louis, MO., USA) No. R41403, HILLIER et al., "yf94c12.s1 Homo sapeins cDNA clone 30225 3', 22 May 1995, see sequence alignment.	1-13

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

20 NOVEMBER 1996

Date of mailing of the international search report

29 NOV 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

GARNETTE D. DRAPER

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/13282

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Database EST-STN on MASPAS search, GenBank at National Library of Medicine, No. S78214, 10 July 1992, MIKI et al., "Disruption of the APC gene by retrotransposal insertion of L1 sequence in a colon cancer", Cancer Research, Vol. 52 No.3, 1992, pages 643-645, see sequence alignment.	1-13
A	EMBL/GENBANK/DDBJ DATA BANKS, No. P41086, EMBL U02603, WOOD, "Putative Succinate Dehydrogenase 15 KD Hydrophobic Protein:", 01 February 1995, see sequence alignment.	1-13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13282

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-13

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US96/13282

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-13, drawn to nucleic acids encoding endokine proteins, a method of making the endokine proteins, and the endokine proteins per se.

Group II, claim 14, drawn to antibodies to the endokine proteins.

Group III, claim 15, drawn to a method of using the antibodies to the endokine proteins in a therapeutic manner.

Group IV, claim 16, drawn to a method of diagnosing for endokine gene expression.

The inventions listed as Groups I, II, III and IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The special technical feature of Group I is directed to nucleic acid and protein products, which are chemically, physically and functionally distinct, and do not require the antibody product of Group III. Furthermore, there is no common special technical feature between the Group I products and the methods of Groups III and IV. Likewise, there is no special technical feature between the method of Group III, for a therapeutic use of the antibody, and the method of Group IV to a diagnostic method of assaying gene expression, because each of these distinct methods require the use of different process steps and elements, and they produce different final outcomes. Therefore, the Groups are not so linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept.